arXiv:math/0108003v1 [math.AG] 1 Aug 2001

Proceedings of Symposia in Pure Mathematics

Field of moduli and field of definition of Galois covers

Stefan Wewers

Abstract. In this paper we investigate the cohomological obstruction for the field of moduli of a G-cover to be a field of definition, in the case of local fields and covers with tame admissible reduction. This applies in particular to p-adic fields where p does not divide the order of the group G. We give examples of G-covers with field of moduli $\mathbb{Q}_p$ that cannot be defined over $\mathbb{Q}_p$, for all primes $p > 5$.

Introduction

In the context of the regular inverse Galois problem, the method of *rigidity* and its generalizations using the braid action and Hurwitz spaces have been very successful. One drawback of these methods is that they apply a priori only to groups with trivial center. To give an example, let G be a finite group and $f : Y \to \mathbb{P}^1$ a G-Galois cover of the projective line defined over $\bar{\mathbb{Q}}$. Assume that f has rational branch points $z_1, \ldots, z_r \in \mathbb{P}^1(\mathbb{Q})$ and that the associated tuple of conjugacy classes $(C_1, \ldots, C_r)$ is rational and rigid. Then $\mathbb{Q}$ is the *field of moduli* of the G-cover, i.e. for each $\sigma \in \mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ the conjugate cover ${}^{\sigma}f$ is isomorphic to f. If, moreover, G has trivial center, then $\mathbb{Q}$ is a *field of definition* of f, i.e. there exists a model $f_{\mathbb{Q}} : Y_{\mathbb{Q}} \to \mathbb{P}^1_{\mathbb{Q}}$ of f over $\mathbb{Q}$. This yields the basic Rigidity Criterion, see e.g. [**17**], [**22**] or [**24**].

However, if the center C of G is not trivial, the field of moduli k_m of a G-cover need not be a field of definition. There is an obstruction $\omega \in H^2(k_m, C)$ such that f can be defined over an extension k/k_m if and only if the restriction of ω to k vanishes. This cohomological approach has been developed in several papers, in the context of Galois covers and in more general situations, see e.g. [**4**], [**5**]. In [**6**] it is used to prove a local-to-global principle: a G-cover with field of moduli $\mathbb{Q}$ is defined over $\mathbb{Q}$ if and only if it can be defined over $\mathbb{Q}_p$, for each prime number p. In [**8**] it is shown that $\omega|_{\mathbb{Q}_p}$ vanishes for each prime number p at which the cover f has good reduction; this includes all “good” primes in the sense of [**2**]. In the present paper we describe a method that allows, in many cases, to explicitly compute the local obstruction $\omega|_{\mathbb{Q}_p}$, at a prime p at which the cover f has bad reduction (but which does not divide the order of $|G|$). We also apply this method in a nontrivial example.

©0000 (copyright holder)

To explain our results in more detail, let k be a field which is henselian with respect to a discrete valuation v, and let $f : Y \to \mathbb{P}^1$ be a G-cover, defined over the algebraic closure of k, with field of moduli k. Our main assumption is that f has *tame admissible* reduction. For instance, this condition holds if the order of G is prime to the residue characteristic of k. In this case, bad reduction is caused by degeneration of the branch locus of the cover modulo v. In particular, the results we present here are interesting only for covers with at least 4 branch points. By standard lifting results, the special fiber of a tame admissible cover, together with certain degeneration data, 'knows everything' about the generic fiber, see e.g. [**26**]. Therefore, one can hope to compute the obstruction for k to be a field of definition purely in terms of the special fiber of f.

Consider the following short exact sequence of cohomology groups:

$$0 \to H^2(k_0, C) \longrightarrow H^2(k, C) \xrightarrow{r_v} H^1(k_0, C(-1)) \to 0, \tag{0.1}$$

where k_0 denotes the residue field of k, see [**23**], Section II.A.2. Suppose $f : Y \to \mathbb{P}^1$ is a G-cover with field of moduli k and $\omega \in H^2(k, C)$ is the associated obstruction for k to be a field of definition. The class $r_v(\omega) \in H^1(k_0, C(-1))$ is called the *residue* of ω at v. If $r_v(\omega) = 0$, then ω, regarded as a class in $H^2(k_0, C)$, represents the obstruction for the special fiber $\bar{f}$ of f to be defined over k_0 (as an admissible G-cover). This happens, for instance, if f has good reduction. On the other hand, if $H^2(k_0, C) = 0$ (for instance, if k_0 is a finite field) then the special fiber $\bar{f}$ has a k_0-model $\bar{f}_{k_0}$, and then $r_v(\omega)$ represents the obstruction for $\bar{f}_{k_0}$ to lift to a k-model of f. These considerations immediately give a new proof of the main result of [**8**].

If the residue field k_0 has characteristic 0, one can describe the degeneration behavior of f purely in terms of the *Hurwitz data* attached to f. A substantial part of the present paper is devoted to showing that, in many cases, this description is sufficient for computing $r_v(\omega)$.

To compute ω over a p-adic field, we use a specialization technique. Given a p-adic field k and a G-cover f with field of moduli k, one can construct (under certain conditions) a G-cover f_t, with field of moduli $K_t := k((t))$, such that $f = f_a$ is a 'specialization' of f_t, for some value $a \in k$ with $v(a) > 0$. Since k has characteristic 0, one can apply the more geometric methods referred to in the preceding paragraph to compute the obstruction ω_t for K_t to be a field of definition of f_t. In this situation, we prove that the obstruction ω corresponding to f can be computed by 'specializing' ω_t to $t = a$. The proof uses the theory of Hurwitz spaces in mixed characteristic, see e.g. [**25**].

Let $\tilde{A}_5$ be the unique nonsplit central extension of A_5 by ± 1. We construct, for each prime number $p > 5$, an $\tilde{A}_5$ cover with field of moduli $\mathbb{Q}_p$ which is not defined over $\mathbb{Q}_p$. To my knowledge, this kind of example is essentially new ([**4**], Example 2.6, for instance, uses only the real numbers).

We should compare this paper to several approaches in the literature to understand the absolute Galois group of $\mathbb{Q}$ acting on the fundamental groups of moduli spaces. [**16**] provides a Hurwitz space context for the Drinfeld-Ihara-Grothendieck relations (that apply to elements of the absolute Galois group; call these DIG relations). The approach was through *tangential base points* . We follow that tradition, though we are not taking the exact same tangential base points. For example, we often use complex conjugate pairs of branch points, while they always used sets of real branch points. Ihara's use of the DIG relations has been primarily to describe the Lie algebra of the absolute Galois group acting through various pronilpotent braid groups, especially on the 3 punctured λ-line. Fried in [**Fri95a**, App. C] proposed Modular Towers, a profinite construction, as suitably like finite representations of the fundamental group to see the DIG relations at a *finite level* . There is an analogy with Ihara in that modular curves are close to considerations about the λ-line, though there is no phenomenon related to them that suggests seeing the DIG relations. Still, modular curves are just one case of Modular Towers.

Fried specifically proposed the Modular Tower attached to A_5 and four 3-cycles as a candidate where one would see a system of Serre obstruction situations from covers of A_5. [**1**, Prop. 8.12] uses a specific case of Modular Tower levels having a tower of these Serre obstructions. Over the reals, there are points, *Harbater-Mumford (HM) reps.* at every level of this Modular Tower where (in our notation) the Serre obstruction vanishes and has value 0, and other points (*near HM reps.*) where it vanishes, but does not have value 0. At levels 1 and beyond, these are the only real points and the real components through them end at cusps of reduced Hurwitz spaces (covering the j-line). These computations are slightly differently than ours, and they give their own approach to Serre's obstruction through direct use of rational points on covers [**1**, Prop. 6.8] . Still, restricting our Prop. 2.17 to the reals is essentially equivalent to an example of theirs.

This suggests there are analogs of p-adic points on these Modular Towers levels that have a similar tangential base point (cusp geometry) analysis to the near HM and HM reps. Further, this geometry should reveal the DIG relations on actual covers, instead of as a Lie algebra relation. While there may be technical difficulties with carrying this out at all levels, the computations of [**1**, Prop. 9.8] at level 1 of this A_5 Modular Tower should be feasible.

The paper is organized in two sections. In Section 1, we recall general results about the field of moduli and the field of definition of Galois covers, from our point of view. Section 2 is concerned with the case of a henselian ground field and contains the main results.

The author would like to thank the organizers of the special semester *Galois Groups and Fundamental Groups* for inviting him and for providing financial support. He would also like to thank the MSRI for its hospitality, and the referee for useful comments. The final version of this paper was written while the author received a grant from the *Deutsche Forschungsgemeinschaft*.

1. General results

In Section 1.1 we recall the necessary definitions. In Section 1.2 we introduce some notation concerning the Hurwitz description of Galois covers and compute the f.o.d.-obstruction in an example (the group is $\mathrm{SL}_2(\ell)$ and the ramification type is $(4,\ell,\ell)$). In Section 1.3 we study in some detail Galois covers over the real numbers.

1.1. The field of moduli condition and the f.o.d.-obstruction. Let G be a finite group and k a field. We denote by k^s a fixed separable closure of k and by $\Gamma_k := \mathrm{Gal}(k^s/k)$ the absolute Galois group of k. Suppose we are given a G-cover

$$f : Y \xrightarrow{G} \mathbb{P}^1_{k^s}$$

of the projective line over k^s; by this we mean that f is a finite Galois cover of smooth projective curves over k^s, with Galois group G. We will always assume that f is *tamely ramified*, and we let $S = \{z_1, \dots, z_r\} \subset \mathbb{P}^1_{k^s}$ be the set of branch points of f. Thus, the G-cover f corresponds to a surjective morphism $\Phi : \pi_1^t(U) \to G$, where $\pi_1^t(U)$ denotes the tame fundamental group of $U := \mathbb{P}^1_{k^s} - S$.

The question we are concerned with is the following: is k a *field of definition* of f, i.e. does there exist a G-cover $f_k : Y_k \to \mathbb{P}^1_k$ defined over k such that $f = f_k \otimes_k k^s$? A necessary condition for this to hold is that the branch locus S is Γ_k-invariant. We will assume this from now on, and let $S_k \subset \mathbb{P}^1_k$ be the closed subset of $\mathbb{P}^1_k$ corresponding to S and $U_k := \mathbb{P}^1_k - S_k$ its complement. By definition, k is a field of definition of f if and only if Φ can be extended to a morphism $\Phi_k : \pi_1^t(U_k) \to G$:

$$\begin{tikzcd}
1 \arrow[r] & \pi_1^t(U) \arrow[r] \arrow[d, "\Phi"] & \pi_1^t(U_k) \arrow[r] \arrow[dl, "\Phi_k"] & \Gamma_k \arrow[r] & 1 \\
 & G & & &
\end{tikzcd}$$

Let us fix a section $s : \Gamma_k \to \pi_1^t(U_k)$ of the natural projection $\pi_1^t(U_k) \to \Gamma_k$, and let Γ_k act on $\pi_1^t(U)$ as follows: ${}^{\sigma}\gamma := s(\sigma)\,\gamma\, s(\sigma)^{-1}$.

Definition 1.1. We say that k is a *field of moduli* of the G-cover $f : Y \to \mathbb{P}^1_{k^s}$ if for all $\sigma \in \Gamma_k$ there exists an element $h_\sigma \in G$ such that

$$\Phi({}^{\sigma}\gamma\,) \;=\; h_\sigma\,\Phi(\gamma)\,h_\sigma^{-1}, \qquad \text{for all } \gamma \in \pi_1^t(U).$$

Note that the field of moduli condition does not depend on the choice of the section s. Obviously, in order for k to be a field of definition of f, it is necessary that k be a field of moduli. However, this is not a sufficient condition, in general. Let C be the center of G and $\bar{G} := G/C$ the quotient. Assume that k is a field of moduli of f, let h_σ, for each $\sigma \in \Gamma_k$, be as in Definition 1.1 and set $\bar{\varphi}(\sigma) := \bar{h}_\sigma$ (the class of h_σ in $\bar{G}$). Clearly, $\bar{\varphi} : \Gamma_k \to \bar{G}$ is well defined and a group homomorphism. It follows that for all $\sigma, \tau \in \Gamma_k$, the element

$$c_{\sigma,\tau} := h_\sigma\, h_\tau\, h_{\sigma\tau}^{-1}$$

lies in the center C, and that $(\sigma,\tau) \mapsto c_{\sigma,\tau}$ is a 2-cocycle (where C is regarded as constant Γ_k-module). Let $\omega \in H^2(k, C)$ be the cohomology class of this cocycle. By definition, ω is the obstruction for the existence of a (weak) solution φ of the central embedding problem $(q, \bar{\varphi})$, see e.g. [**17**, IV.6.1] (here $q : G \to \bar{G}$ is the natural map).

$$\begin{tikzcd}
 & & & \Gamma_k \arrow[dl, "\varphi"'] \arrow[d, "\bar{\varphi}"] & \\
1 \arrow[r] & C \arrow[r] & G \arrow[r, "q"] & \bar{G} \arrow[r] & 1
\end{tikzcd} \tag{1.1}$$

A formal verification shows (see e.g. [**5**]):

Proposition 1.2. *The class ω is independent of the choice of the section $s : \Gamma_k \to \pi_1^t(U_k)$. Moreover, $\omega = 0$ if and only if k is a field of definition of f.*

We call ω the *f.o.d.-obstruction* for the G-cover f, relative to k ('f.o.d.' stands for 'field of definition'). The rest of the paper is concerned with studying and, if possible, computing ω, in various situations. First we make some general remarks.

Remark 1.3. (i) The formation of ω is functorial in k. More precisely, let K/k be a field extension, $f_{K^s} : Y \otimes_{k^s} K^s \to \mathbb{P}^1_{K^s}$ the base change of f to K^s (we embed k^s in K^s) and $\omega|_K$ the image of ω under the restriction homomorphism $H^2(k,C) \to H^2(K,C)$; then $\omega|_K$ is the f.o.d.-obstruction for f_{K^s}.

(ii) A G-cover $f : Y \to \mathbb{P}^1_{k^s}$ with r branch points corresponds to a k^s-point $[f] : \operatorname{Spec} k^s \to \mathcal{H}_r(G)$ on a certain moduli scheme $\mathcal{H}_r(G)$, called the *Hurwitz space*. The field k is a field of moduli for f if and only if $[f]$ is k-rational, i.e. factors through a morphism $\operatorname{Spec} k \to \mathcal{H}_r(G)$. See [**10**] and [**25**].

(iii) There exists a cohomology class $\tilde\omega \in H^2_{\text{ét}}(\mathcal{H}_r(G), C)$ which specializes to ω, i.e. $\omega = [f]^*\tilde\omega$. The class $\tilde\omega$ represents the obstruction for the existence of a global versal G-cover over $\mathcal{H}_r(G)$, see [**25**] or [**7**].

For a study of f.o.d.-obstructions for more general (e.g. non-Galois) covers, see [**5**] and [**7**].

1.2. The Hurwitz description. We let k, $S = \{z_1, \dots, z_r\}$, $U := \mathbb{P}^1_{k^s} - S$ and U_k be as before. We assume that k has characteristic 0; we may therefore write $\pi_1(U)$ instead of $\pi_1^t(U)$. Let us choose a k-rational *base point* z_0 on U. The point z_0 will either be a k-rational point $\operatorname{Spec} k \to U_k$ or a "tangential base point" $\operatorname{Spec} k((z)) \to U_k$, see [**15**]. In both cases, we obtain a section $s : \Gamma_k \to \pi_1(U_k)$ (unique up to an inner automorphism of $\pi_1(U)$) and hence an action of Γ_k on $\pi_1(U)$. We will write $\pi_1(U, z_0)$ to denote the profinite group $\pi_1(U)$ together with this Γ_k-action.

We denote by $\hat{\mathbb{Z}}(1) := \varprojlim_n \boldsymbol{\mu}_n$ the Tate module of $\mathbb{G}_m$. Let

$$\Pi \ := \ <\gamma_1, \dots, \gamma_r \mid \prod_i \gamma_i = 1> \tag{1.2}$$

be the free profinite group with generators $\gamma_1, \dots, \gamma_r$, subject to the usual product one relation.

Definition 1.4. An isomorphism $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ of profinite groups is a *presentation* of $\pi_1(U, z_0)$ if it has these properties:

(i) $\rho(\gamma_i)$ generates an inertia subgroup $I_{z_i} \subset \pi_1(U, z_0)$ corresponding to a point $z_i \in S$, and

(ii) under the natural identification $I_{z_i} \cong \hat{\mathbb{Z}}(1)$, all the $\rho(\gamma_i)$ correspond to the same element in $\hat{\mathbb{Z}}(1)$.

It is a well known fact that such a presentation ρ always exists; let us choose one. To simplify the notation, we will usually identify γ_i with $\rho(\gamma_i)$. The *branch cycle argument* states that for all $\sigma \in \Gamma_k$ there exist elements $\beta_1, \dots, \beta_r \in \pi_1(U, z_0)$ such that

$$^{\sigma}\gamma_i \ = \ \beta_i \, \gamma_{\sigma(i)}^{\chi(\sigma)} \, \beta_i^{-1}. \tag{1.3}$$

Here $\chi : \Gamma_k \to \hat{\mathbb{Z}}^\times$ is the cyclotomic character and $\sigma(i)$ is defined by $\sigma(z_i) = z_{\sigma(i)}$.

Let $f : Y \to \mathbb{P}^1_{k^s}$ be a G-cover with branch locus S, corresponding to a homomorphism $\Phi : \pi_1(U, z_0) \to G$. Setting $g_i := \Phi(\gamma_i)$, we obtain an r-tuple $\mathbf{g} = (g_1, \dots, g_r)$ of generators of G such that $\prod_i g_i = 1$. We write $\mathrm{Ni}_r(G)$ for the set of all such $\mathbf{g}$ and $\mathrm{Ni}_r^{\mathrm{in}}(G) := \mathrm{Ni}_r(G)/\operatorname{Inn}(G)$ for the quotient of this set under the action of inner automorphisms of G, compare [**10**]. We obtain a bijection between $\mathrm{Ni}_r^{\mathrm{in}}(G)$ (the set of *Nielsen classes* of length r) and the set of isomorphism classes of G-covers with branch locus S. We say that the G-cover f has *Hurwitz description* $[\mathbf{g}] \in \mathrm{Ni}_r^{\mathrm{in}}(G)$ (with respect to the presentation ρ).

The Γ_k-action on $\pi_1(U,z_0)$ induces a Γ_k-action on $\mathrm{Hom}(\pi_1(U,z_0),G)$, and therefore on $\mathrm{Ni}_r(G)$. Our convention is to let Γ_k act on $\mathrm{Ni}_r(G)$ from the *right*, i.e. we define

$$\mathbf{g}^\sigma := (\tilde g_1,\dots,\tilde g_r), \quad \text{with} \quad \tilde g_i := \Phi(\sigma\,\gamma_i\,\sigma^{-1}).$$

We remark that this is not a standard convention in the literature. The branch cycle argument (1.3) becomes

$$\tilde g_i = b_i\, g_{\sigma(i)}^{\chi(\sigma)}\, b_i^{-1},$$

with $b_i := \Phi(\beta_i)$. The Γ_k-action on $\mathrm{Ni}_r(G)$ induces a Γ_k-action on $\mathrm{Ni}_r^{\mathrm{in}}(G)$. The following Proposition merely rephrases the definitions of Section 1.1.

Proposition 1.5. *Let $f : Y \to \mathbb{P}^1_{k^s}$ be a G-cover with branch locus S and Hurwitz description $\mathbf{g}$ (with respect to a presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U,z_0)$). Then*

(i) *k is a field of moduli of f if and only if $[\mathbf{g}]^\sigma = [\mathbf{g}]$ for all $\sigma \in \Gamma_k$.*

(ii) *If (i) holds then there is a unique homomorphism $\bar\varphi : \Gamma_k \to \bar G$ such that*

$$\mathbf{g}^\sigma = h_\sigma\,\mathbf{g}\,h_\sigma^{-1},$$

where $h_\sigma \in G$ is a lift of $\bar\varphi(\sigma) \in \bar G$, for all $\sigma \in \Gamma_k$.

(iii) *k is a field of definition of f if and only if it is a field of moduli and the homomorphism $\bar\varphi$ in (ii) lifts to a homomorphism $\varphi : \Gamma_k \to G$.*

The formula $\mathbf{g}^\sigma = h_\sigma\,\mathbf{g}\,h_\sigma^{-1}$ may seem counterintuitive at first. But we have to keep in mind that the homomorphism $\bar\varphi$ depends on $\mathbf{g}$, and not only on its Nielsen class $[\mathbf{g}]$. In more geometric terms, $\mathbf{g}$ corresponds to a *pointed G-cover* $f : (Y,y_0) \to (\mathbb{P}^1,z_0)$. If k is a field of definition of f and $\varphi : \Gamma_k \to G$ is as in Proposition 1.5 (iii), then there exists a unique model $f_k : Y \to \mathbb{P}^1_k$ of f such that the fiber $f^{-1}(z_0)$, as a G-torsor over k, corresponds to $\varphi \in H^1(k,G)$.

Example 1.6. Let $r := 3$, $k := \mathbb{Q}$, and let ℓ be an odd prime. We set $\ell^* := (-1)^{(\ell-1)/2}\ell$, $S := \{0,\sqrt{\ell^*},-\sqrt{\ell^*}\} \subset \mathbb{P}^1_{\bar{\mathbb{Q}}}$ and $U := \mathbb{P}^1_{\bar{\mathbb{Q}}} - S$. We identify $\mathbb{Q}(z)$ with the function field of $U_{\mathbb{Q}} := \mathbb{P}^1_{\mathbb{Q}} - S$ and let $z_0 : \mathrm{Spec}\,\mathbb{Q}((z)) \to U_{\mathbb{Q}}$ be the tangential base point at 0, with parameter z.

It is clear from [**15**] that there exists a presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U,z_0)$ such that $\gamma_1 \in \pi_1(U,z_0)$ corresponds to the closed path $t \mapsto e^{2\pi i t}$ (for $0 \le t \le 1$). Then

$$(1.4) \qquad {}^\sigma\gamma_1 = \gamma_1^{\chi(\sigma)}, \qquad {}^\sigma\gamma_2 \sim \gamma_{\sigma(2)}, \qquad {}^\sigma\gamma_3 \sim \gamma_{\sigma(3)},$$

for all $\sigma \in \Gamma_{\mathbb{Q}}$. Here σ fixes the indices 2 and 3 if ${}^\sigma\sqrt{\ell^*} = \sqrt{\ell^*}$ and permutes them if ${}^\sigma\sqrt{\ell^*} = -\sqrt{\ell^*}$.

From now on we assume that $\ell \not\equiv \pm 1 \pmod 8$, and we set $G := \mathrm{SL}_2(\ell)$. Let ℓA, ℓB be the two conjugacy classes of G containing elements of order ℓ and $4A$ the unique conjugacy class containing elements of order 4. The classes ℓA and ℓB are conjugate over the quadratic extension $\mathbb{Q}(\sqrt{\ell^*})/\mathbb{Q}$. By [**24**] I.3.3.6, the triple $(4A,\ell A,\ell B)$ is *rigid*, i.e. there exists exactly one class $[g_1,g_2,g_3] \in \mathrm{Ni}_3^{\mathrm{in}}(G)$ such that $g_1 \in 4A$, $g_2 \in \ell A$ and $g_3 \in \ell B$. Let $f : Y \to \mathbb{P}^1_{\bar{\mathbb{Q}}}$ be the G-cover with Hurwitz description $[\mathbf{g}] = [g_1,g_2,g_2]$ (with respect to ρ). By Rigidity and (1.4), $[\mathbf{g}]^\sigma = [\mathbf{g}]$ for all $\sigma \in \Gamma_{\mathbb{Q}}$. Therefore, $\mathbb{Q}$ is a field of moduli of f.

The center C of G consists of the two diagonal matrices $\pm I$; we identify it with $\{\pm 1\}$. Let $\omega \in H^2(\mathbb{Q},\pm 1)$ be the f.o.d.-obstruction of f. We may identify $H^2(\mathbb{Q},\pm 1)$ with $\mathrm{Br}_2(\mathbb{Q})$, the 2-torsion of the Brauer group of $\mathbb{Q}$.

Proposition 1.7. *We have $\omega = (-1,-1)$, i.e. ω is represented (as an element of $\mathrm{Br}_2(\mathbb{Q})$) by the quaternion algebra $\mathbb{Q}[i,j \mid i^2 = -1, j^2 = -1, ij = -ji]$. In particular, $\mathbb{Q}$ is not a field of definition of f.*

It follows that the regular $\mathrm{PSL}_2(\ell)$-extensions of $\mathbb{Q}(z)$ with ramification of type $(2,\ell,\ell)$ (see e.g. [**22**], Section 8.3.3) do not lift to an $\mathrm{SL}_2(\ell)$-extension. We will see in Section 1.3 that in fact no $\mathrm{PSL}_2(\ell)$-extension with three branch points lifts to an $\mathrm{SL}_2(\ell)$-extension.

Proof. The class $\omega \in H^2(\mathbb{Q},\pm 1)$ is the obstruction for lifting the homomorphism $\bar\varphi : \Gamma_{\mathbb{Q}} \to \bar G = \mathrm{PSL}_2(\ell)$ to a homomorphism $\varphi : \Gamma_{\mathbb{Q}} \to G = \mathrm{SL}_2(\ell)$. By the definition of $\bar\varphi$ and by Equation (1.4), we have (by abuse of notation)

$$\bar\varphi(\sigma)\, g_1\, \bar\varphi(\sigma)^{-1} \;=\; g_1^{\chi(\sigma)}, \qquad \text{for all } \sigma \in \Gamma_{\mathbb{Q}}. \tag{1.5}$$

In particular, the image of $\bar\varphi$ is contained in $\bar N := N/C$, where N is the normalizer in G of the cyclic subgroup generated by g_1. It is shown e.g. in [**14**], Abschnitt II.8, that $\bar N$ is a dihedral group of order $4n$, where $n := (\ell-1)/4$ if $\ell \equiv 1 \pmod 4$ and $n := (\ell+1)/4$ otherwise. Let $\bar H \lhd \bar N$ be the unique cyclic normal subgroup of order n. By our assumption on ℓ, n is odd. Therefore, there exists a unique cyclic normal subgroup $H \lhd N$ of order n mapping onto $\bar H$. Since $\bar N/\bar H \cong \mathbb{Z}/2\times\mathbb{Z}/2$ and every involution in $\bar G$ lifts to an element of order 4 in G, N/H is isomorphic to the quaternion group Q_8. It is easy to check that in the following diagram

$$\begin{tikzcd} \{\pm 1\} \arrow[r] \arrow[d] & N \arrow[r] \arrow[d] & \bar N \arrow[d] \\ \{\pm 1\} \arrow[r] & N/H \cong Q_8 \arrow[r] & N/CH \cong \mathbb{Z}/2\times\mathbb{Z}/2 \end{tikzcd} \tag{1.6}$$

the group extension in the upper row is the pullback of the lower row with respect to the projection $\bar N \to N/CH$. Let $\bar\psi : \Gamma_{\mathbb{Q}} \to N/CH$ be the composition of $\bar\varphi$ with the projection $\bar N \to N/CH$. It follows from [**3**], Exercise IV.3.1, that $\omega = \bar\psi^*\eta$, where $\eta \in H^2(N/CH, C)$ is the cohomology class corresponding to the lower row of (1.6). In other words, ω is the obstruction for lifting $\bar\psi$ to a homomorphism $\psi : \Gamma_{\mathbb{Q}} \to N/H \cong Q_8$. Therefore, by a result of Witt (see [**11**], (7.7) (ii)),

$$\omega = (-1,a) + (-1,b) + (a,b) \in \mathrm{Br}_2(\mathbb{Q}),$$

where $E = \mathbb{Q}[x,y \mid x^2 = a,\, y^2 = b]$, $a,b \in \mathbb{Q}$, is the étale $\mathbb{Q}$-algebra corresponding to $\bar\psi$. Let $\bar H' \lhd \bar N$ be the maximal cyclic normal subgroup, of order $2n$. By (1.5), the composition of $\bar\psi$ with $N/CH \to \bar N/\bar H' \cong \mathbb{Z}/2 \cong (\mathbb{Z}/4)^\times$ equals the cyclotomic character modulo 4. Hence we may choose $a = -1$, and $\omega = (-1,-1)$ follows. □

1.3. *G*-covers over the real numbers.

In this section we discuss the case $k = \mathbb{R}$. Since the action of $\Gamma_{\mathbb{R}}$ on $\pi_1(U,z_0)$ is known, we get some easy results and examples which illustrate several key concepts of this paper. For more general results on related questions, see [**1**], in particular §6.

Let $f : Y \to \mathbb{P}^1_{\mathbb{C}}$ be a G-cover defined over the complex numbers, and assume that the branch locus S of f is defined over $\mathbb{R}$. For simplicity, we will also assume that $\infty \notin S$. We can write $S = \{z_1,\ldots,z_r\}$, with $\bar z_i = z_{2s-i+1}$ for $i = 1,\ldots,s$, and $z_{2s+j} \in \mathbb{R}$ for $j = 1,\ldots,r-2s$. For $z,w \in \mathbb{C}$, we write $z \succ w$ if either $\operatorname{Im} z > \operatorname{Im} w$ or $\operatorname{Im} z = \operatorname{Im} w$ and $\operatorname{Re} z > \operatorname{Re} w$. In this notation, we may assume that $z_1 \succ \ldots \succ z_s$ and $z_{2s+1} < \ldots < z_r$. Let $U := \mathbb{P}^1_{\mathbb{C}} - S$ and choose a real base point

$z_0 < z_{2s+1}$. By [**17**, Section I.1.1], there exists a presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ with the following properties: (i) γ_i is represented by a simple loop based at z_0 winding counterclockwise around z_i, (ii) these loops are pairwise disjoint and (iii) the action of complex conjugation $\kappa \in \Gamma_{\mathbb{R}}$ is given by

$$ {}^{\kappa}\gamma_i \;=\; \gamma_{2s-i+1}^{-1}, \qquad \text{for } i = 1, \dots, 2s, \tag{1.7}$$

and

$$ {}^{\kappa}\gamma_{2s+j} \;=\; \beta_j\, \gamma_{2s+j}^{-1}\, \beta_j^{-1}, \qquad \text{for } j = 1, \dots, r-2s, \tag{1.8}$$

where $\beta_j := \gamma_{2s+1} \cdot \ldots \cdot \gamma_{2s+j-1}$ (see [**17**, Fig. 1.2]). We say that ρ is a *normalized presentation* of $\pi_1(U, z_0)$.

PROPOSITION 1.8. *Let $\mathbf{g} = (g_1, \dots, g_r)$ be the Hurwitz description for the G-cover $f : Y \to \mathbb{P}^1_{\mathbb{C}}$, with respect to a normalized presentation ρ, as above. Then*

(i) *$\mathbb{R}$ is a field of moduli for f if and only if there exists $b \in G$ such that*

$$ b\, g_i\, b^{-1} \;=\; g_{2s-i+1}^{-1} $$

for $i = 1, \dots, 2s$, and

$$ b\, g_{2s+j}\, b^{-1} \;=\; h_j\, g_{2s+j}^{-1}\, h_j^{-1} $$

for $j = 1, \dots, r-2s$, with $h_j := g_{2s+1} \cdot \ldots \cdot g_{2s+j-1}$.

(ii) *$\mathbb{R}$ is a field of definition of f if and only there exists b as in* (i) *such that $b^2 = 1$.*

PROOF. Follows from Proposition 1.5 and equations (1.7) and (1.8). □

For certain groups G, the condition in Proposition 1.8 (ii) is rather restrictive, as shown by the next theorem.

THEOREM 1.9. *Let G be a finite group. Assume that all elements of G of order 2 lie in the center of G. Let $f : Y \to \mathbb{P}^1_{\mathbb{C}}$ be a G-cover, with branch locus S defined over $\mathbb{R}$, and let $\mathbf{g}$ be a Hurwitz description of f with respect to a normalized presentation $\rho : \Pi \to \pi_1(U, z_0)$ (see above). Then $\mathbb{R}$ is a field of definition of f if and only if*

$$ \mathbf{g} = (\,g_1, \dots, g_s, g_s^{-1}, \dots, g_1^{-1},\, c_1, \dots, c_{r-2s}), \tag{1.9}$$

with $c_j^2 = 1$ and $\prod_j c_j = 1$.

The quaternion group Q_8 and $\mathrm{SL}_2(\ell^n)$, for $\ell \neq 2$ verify the conditions of the theorem. Elements of order 2 in A_n are products of $2s$ disjoint 2-cycles. They lift to have order 4 in the 2-fold coverings $\tilde{A}_n$ of A_n exactly when s is odd [**1**, Prop. 5.8]. So this also holds for $\tilde{A}_n$ if $4 \leq n \leq 7$.

PROOF. If $\mathbf{g}$ is as in (1.9), then ${}^{\kappa}\mathbf{g} = \mathbf{g}$, so f can be defined over $\mathbb{R}$. Conversely, if f can be defined over $\mathbb{R}$ then there exists $b \in G$ of order 2 such that $b\, g_i\, b^{-1} = g_{2s-i+1}^{-1}$ and $b\, g_{2s+j}\, b^{-1} = h_j\, g_{2s+j}^{-1}\, h_j^{-1}$, by Proposition 1.8 (ii). By the hypothesis of the theorem, b is a central element of G. Therefore, $g_i = g_{2s-i+1}^{-1}$, for $i = 1, \dots, s$. Moreover, $g_{2s+1} = g_{2s+1}^{-1}$, so $c_1 := g_{2s+1}$ has order 2 and hence lies in the center of G. Continuing by induction, we find that $c_j := g_{2s+j}$ is central of order 2, for $j = 1, \dots, r-2s$. This proves the theorem. □



FIGURE 1.

EXAMPLE 1.10. Let $G := \tilde{A}_5$ be the nonsplit central extension of A_5 by ± 1. Let $g_1, g_2 \in G$ be the unique lifts of order 3 of $\bar{g}_1 := (1\,2\,3)$, $\bar{g}_2 := (3\,4\,5) \in \bar{G} = A_5$. Set $\mathbf{g} := (\,g_1,\, g_2,\, g_2^{-1}, g_1^{-1}) \in \mathrm{Ni}_4(G)$. For $t \in \mathbb{C}$, set

$$S_t \;:=\; \{\; z \in \mathbb{C} \mid z^4 + (2t-2)z^2 + t^2 + 2t + 1 = 0 \;\} \;=\; \{\; \pm 1 \pm \sqrt{-t}\;\}$$

and $U_t := \mathbb{P}^1_{\mathbb{C}} - S_t$. Let us fix a value $t_0 \in \mathbb{R}$ such that $0 < t_0 < 1$, and let ρ_{t_0} be the normalized presentation of $\pi_1(U_{t_0}, 0)$, as shown on the left hand side of Figure 1. We let $f_{t_0} : Y_{t_0} \to \mathbb{P}^1_{\mathbb{C}}$ be the G-cover with Hurwitz description $\mathbf{g}$, with respect to ρ_{t_0}. By Theorem 1.9, f_{t_0} can be defined over $\mathbb{R}$.

Let $\theta : [0,1] \to \mathbb{C}$ be the path $s \mapsto \theta_s := t_0\, e^{\pi i s}$, and let $f_{-t_0} : Y_{-t_0} \to \mathbb{P}^1_{\mathbb{C}}$ be the $\tilde{A}_5$-cover with branch locus S_{-t_0} obtained from f_{t_0} by "analytic continuation of the branch locus along $s \mapsto S_{\theta_s}$" (see e.g. [**24**, Section II.10.1] for precise definitions). Thus, f_{-t_0} has branch locus $S_{-t_0} = \{\pm 1 \pm \sqrt{t_0}\}$, and Hurwitz description $\mathbf{g}$ with respect to the presentation ρ_{-t_0}, as shown on the right hand side of Figure 1. Note that the presentation ρ_{-t_0} is not normalized. From Figure 1, we see that the action of complex conjugation on $\pi_1(U_{-t_0}, 0)$, with respect to ρ_{-t_0}, is given by

$$^{\kappa}\gamma_1 = \gamma_1^{-1}, \quad {}^{\kappa}\gamma_2 = \gamma_3^{-1}\,\gamma_2^{-1}\,\gamma_3, \quad {}^{\kappa}\gamma_3 = \gamma_3^{-1}, \quad {}^{\kappa}\gamma_4 = \gamma_1^{-1}\,\gamma_4^{-1}\,\gamma_1. \tag{1.10}$$

A short computation reveals how this new action affects the tuple $\mathbf{g}$:

$$\mathbf{g}^{\kappa} \;=\; (\,g_1^{-1}, g_2^{-1}, \,g_2,\, g_1\,) \;=\; b\,\mathbf{g}\,b^{-1}, \tag{1.11}$$

where $b \in \tilde{A}_5$ is a lift of $(1\,2)(4\,5) \in \bar{G} = A_5$. Therefore, $\mathbb{R}$ is a field of moduli of f_{-t_0}. On the other hand, any element $b \in \tilde{A}_5$ such that (1.11) holds has order 4 (there are exactly two of them), so $\mathbb{R}$ is not a field of definition of f_{-t_0}. In other words: for $t \in (-1,1) - \{0\}$, let $\omega_t \in H^2(\mathbb{R}, \{\pm 1\})$ be the f.o.d.-obstruction for f_t. Then $\omega = 0$ for $t > 0$ and $\omega = (-1,-1) \neq 0$ for $t < 0$.

Example 1.10 illustrates a key point of this paper: the obstruction ω_t, as a function of $t \in (-1,1) \subset \mathbb{R}$, has a 'jump' at $t = 0$. In the next section we will give the following algebraic interpretation for this phenomenon. Let $\tilde{f} : \tilde{Y} \to \mathbb{P}^1_{k^s}$ be the G-cover over k^s, with $k = \mathbb{R}((t))$, corresponding to the family f_t in an infinitesimal neighborhood of $t = 0$. Then k is a field of moduli for $\tilde{f}$, and the corresponding f.o.d.-obstruction $\omega \in H^2(k, \{\pm 1\})$ has a *pole* at $t = 0$, i.e. the *residue* $r_v(\omega) \in H^1(\mathbb{R}, \{\pm 1\})$ of ω at the valuation v of k does not vanish.

2. Galois covers over henselian fields

In Section 2.1 we recall the definition of the residue map and we state a theorem that says that the residue of the f.o.d.-obstruction of a G-cover vanishes if the cover has good reduction. In Section 2.2 we define *compatible automorphisms* of the free profinite group Π, with respect to an *ordered tree*. This is preparatory work for Section 2.3, where we state some results on the Galois action on $\pi_1(U)$. Here we restrict our attention to henselian fields with residue characteristic 0. In Section 2.4 we explicitly compute the f.o.d.-obstruction for some $\tilde{A}_5$-covers over finite extensions of $\mathbb{Q}((t))$. In Section 2.5 we explain how to 'specialize' our previous results to p-adic fields.

2.1. The residue map. Throughout this section, we assume that k has characteristic 0 and is henselian with respect to a discrete valuation v. We denote the residue field of v by k_0 and assume that k_0 is perfect. We write Γ_k^t for the maximal tame quotient of Γ_k and $I^t \lhd \Gamma_k^t$ for its tame inertia subgroup. We choose a section for the natural map $\Gamma_k^t \to \Gamma_{k_0}$ and identify Γ_k^t with $I^t \rtimes \Gamma_{k_0}$. We remark that, as a Γ_{k_0}-module, I^t is canonically isomorphic to $\hat{\mathbb{Z}}'(1) := \varprojlim_n \mu_n(k_0^s)$, where n runs over the integers prime to the characteristic of k_0.

We are given a G-cover $f : Y \to \mathbb{P}^1_{k^s}$ with branch locus S and field of moduli k. Let $\omega \in H^2(k, C)$ be the f.o.d.-obstruction for f. We choose a k-rational branch point z_0 on $U := \mathbb{P}^1_{k^s} - S$. As explained in Sections 1.1 and 1.2, we obtain a certain homomorphism $\bar{\varphi} : \Gamma_k \to \bar{G}$, and ω is the cohomological obstruction for lifting $\bar{\varphi}$ to a homomorphism $\varphi : \Gamma_k \to G$. We assume that the following tameness condition holds.

Condition 2.1.

(i) The order of C is prime to the characteristic of k_0.
(ii) The homomorphism $\bar{\varphi} : \Gamma_k \to \bar{G}$ is at most tamely ramified at v.

We may therefore regard $\bar{\varphi}$ as a homomorphism $\Gamma_k^t \to \bar{G}$ and ω as the obstruction for lifting $\bar{\varphi}$ to a homomorphism $\varphi : \Gamma_k^t \to G$ (Condition 2.1 (i) implies that there is a natural isomorphism $H^2(k, C) \cong H^2(\Gamma_k^t, C)$).

According to [**23**], Section II.A.2, and Condition 2.1 (i), we have a short exact sequence

$$0 \to H^2(k_0, C) \longrightarrow H^2(k, C) \xrightarrow{r_v} H^1(k_0, C(-1)) \to 0 \tag{2.1}$$

(with $C(-1) := \mathrm{Hom}(I^t, C)$). This sequence can be deduced from the Hochschild–Serre spectral sequence for the group extension determined by the inclusion $I \hookrightarrow \Gamma_k$. The map r_v is called the *residue map*, and $r_v(\omega)$ is called the residue of ω at v. If $r_v(\omega) = 0$ we say that ω is *regular* at v; otherwise, ω is said to have a *pole* at v. If ω is regular at v, we can identify ω with a class in $H^2(k_0, C)$; we denote this class by ω_0 and refer to it as the *value* of ω at v.

Proposition 2.2 below gives a description of $r_v(\omega)$ in terms of $\bar{\varphi}$. To state it, we need some more notation. Let us choose a topological generator q_0 of the inertia group I^t. To simplify the notation, we will identify $C(-1) = \mathrm{Hom}(I^t, C)$ with C (as abelian groups), via $\eta \mapsto \eta(q_0)$. We let Γ_{k_0} act on G as follows:

$$^{\sigma}g \;:=\; \bar{\varphi}(\sigma)\, g\, \bar{\varphi}(\sigma)^{-1}, \qquad \sigma \in \Gamma_{k_0} \subset \Gamma_k^t. \tag{2.2}$$

Finally, let us choose an element $a_0 \in G$ lifting $\bar{\varphi}(q_0) \in \bar{G}$. For all $\sigma \in \Gamma_{k_0}$, we find that

$$c_\sigma := a_0\, ({}^{\sigma}a_0)^{-\chi(\sigma)^{-1}} = a_0\, (({}^{\sigma}a_0)^{\chi(\sigma)^{-1}})^{-1} \tag{2.3}$$

is an element of C.

Proposition 2.2. (i) *The map $\sigma \mapsto c_\sigma$ is a 1-cocycle with values in $C \cong C(-1)$. Its cohomology class equals the residue $r_v(\omega)$.*

(ii) *If* $r_v(\omega)=0$*, then* $\omega_0\in H^2(k_0,C)$ *is the obstruction for lifting the homomorphism* $\bar\varphi_0:=\bar\varphi|_{\Gamma_{k_0}}$ *to* G.

Proof. For each $\sigma\in\Gamma_{k_0}$, let us choose a lift $b_\sigma\in G$ lifting $\bar\varphi(\sigma)$. We may assume that $\sigma\mapsto b_\sigma$ is continuous. Every element $\tau\in\Gamma_k^t$ can be written in a unique way as $\tau=\sigma\,q_0^i$, with $\sigma\in\Gamma_{k_0}$ and $i\in(\hat{\mathbb{Z}}')^\times$. Therefore,

$$\varphi'(\sigma\,q_0^i):=b_\sigma\,a_0^i \tag{2.4}$$

defines a continuous, set-theoretic lift $\varphi':\Gamma_k^t\to G$ of $\bar\varphi$. By definition, the class $\omega\in H^2(k,C)$ is represented by the 2-cocycle

$$\omega_{\tau_1,\tau_2}:=\varphi'(\tau_1)\,\varphi'(\tau_2)\,\varphi'(\tau_1\tau_2)^{-1},\qquad \tau_1,\tau_2\in\Gamma_k^t. \tag{2.5}$$

Using (2.4), one checks that ω_{τ_1,τ_2} is normalized (i.e. equals 1 when one of the τ_j equals 1) and depends only on the class of τ_2 in $\Gamma_{k_0}=\Gamma_k^t/I^t$. By [**23**], Section II.A.1, the residue $r_v(\omega)\in H^1(k_0,\mathrm{Hom}(I^t,C))$ is represented by the 1-cocycle

$$\sigma\;\longmapsto\;(q\mapsto\omega_{q,\sigma}),\qquad \sigma\in\Gamma_{k_0},\ q\in I^t. \tag{2.6}$$

Using (2.4), (2.5) and the identification $C\cong C(-1)=\mathrm{Hom}(I^t,C)$, (2.6) becomes

$$\begin{aligned}\sigma\;\longmapsto\;\omega_{q_0,\sigma}&=\varphi'(q_0)\,\varphi'(\sigma)\,\varphi'(q_0\,\sigma)^{-1}\\&=\varphi'(q_0)\,\varphi'(\sigma)\,\varphi'(\sigma\,q_0^{\chi(\sigma)^{-1}})^{-1}\\&=a_0\,({}^{\sigma}a_0)^{-\chi(\sigma)^{-1}}\;=\;c_\sigma.\end{aligned}$$

This proves the first part of the proposition. The second part is obvious from the definition of the sequence (2.1). □

Corollary 2.3. *Assume that the order of* $\bar\varphi(q_0)\in\bar G$ *is prime to the order of* C*. Then* $r_v(\omega)=0$.

Proof. It follows from our assumption that there exists a lift $a_0\in G$ of $\bar\varphi(q_0)$ such that $\mathrm{ord}\,a_0=\mathrm{ord}\,\bar\varphi(q_0)$ is prime to the order of C. By the definition of c_σ we have

$${}^{\sigma}a_0=(c_\sigma^{-1}\,a_0)^{\chi(\sigma)},$$

for $\sigma\in\Gamma_{k_0}$. Choose an integer n such that $n\equiv1\pmod{\mathrm{ord}\,a_0}$ and $n\equiv0\pmod{\mathrm{ord}\,C}$. Then

$${}^{\sigma}a_0={}^{\sigma}(a_0^n)=(c_\sigma^{-n}\,a_0^n)^{\chi(\sigma)}=a_0^{\chi(\sigma)}.$$

It follows that $c_\sigma=1$, for all $\sigma\in\Gamma_{k_0}$. □

Let us denote by R the ring of integers of k^s. Let $S_R\subset\mathbb{P}^1_R$ be the closure of S in $\mathbb{P}^1_R$. We say that the G-cover $f:Y\to\mathbb{P}^1_{k^s}$ has *good reduction* if (i) S_R is étale over $\mathrm{Spec}\,R$ and (ii) f extends to a finite morphism $f_R:Y_R\to\mathbb{P}^1_R$ which is tamely ramified along S_R and étale everywhere else. If f has good reduction, then $\bar f:\bar Y:=Y_R\otimes k_0^s\to\mathbb{P}^1_{k_0^s}$ is a G-cover, with branch locus $\bar S:=S_R\otimes k_0^s$.

Theorem 2.4. *Assume that* f *has good reduction at* v*. Then* $r_v(\omega)=0$*. Moreover, the value* $\omega_0\in H^2(k_0,C)$ *of* ω *at* v *is the f.o.d.-obstruction for the* G*-cover* $\bar f$ *(for which* k_0 *is a field of moduli).*

This theorem can be seen as a consequence of Remark 1.3 (iii). Let us give a proof which does not rely on the results of [**25**].

Proof. Let $\bar{U} := \mathbb{P}^1_{k_0^s} - \bar{S}$ and $\lambda : \pi_1(U, z_0) \to \pi_1^t(\bar{U}, \bar{z}_0)$ the specialization morphism, defined in [**13**]. Since f is assumed to have good reduction, $\Phi : \pi_1(U, z_0) \to G$ factors through λ and the resulting morphism $\bar{\Phi} : \pi_1^t(\bar{U}, \bar{z}_0) \to G$ corresponds to the reduction $\bar{f}$. Because λ is Γ_k-equivariant and the inertia group $I \lhd \Gamma_k$ acts trivially on $\pi_1^t(\bar{U}, \bar{z})$, the homomorphism $\bar{\varphi} : \Gamma_k \to \bar{G}$ is unramified, i.e. corresponds to a homomorphism $\bar{\varphi}_{k_0} : \Gamma_{k_0} \to \bar{G}$. Thus, Proposition 2.2 implies that $r_v(\omega) = 0$ and that ω_0 is the obstruction for lifting $\bar{\varphi}_{k_0}$ to G, proving the theorem. □

Theorem 2.4 above implies Theorem 3.1 of [**8**]. Namely, assume that the characteristic of k_0 does not divide the order of G and that S_R is étale over $\mathrm{Spec}\, R$. Then f has good reduction, by [**2**], and hence $r_v(\omega) = 0$. If we assume in addition that $\mathrm{cd}\, k_0 \leq 1$ (e.g. if k_0 is finite) then $\omega = 0$, because the sequence (2.1) is exact.

2.2. Ordered trees and compatible automorphisms. Let $T = (V, E)$ be a *tree*, i.e. a finite and simply connected graph. We choose a distinguished vertex $v_0 \in V(T)$ and call it the *root* of T. The choice of v_0 induces a natural orientation on T; we represent an edge $e \in E(T)$ as an ordered pair $e = (v_1, v_2)$ of vertices such that v_1 is closer to v_0 than v_2. For $v \in V(T)$ let $p_v = (v_0, \ldots, v)$ be the shortest path leading from v_0 to v. We define T_v as the subtree of T which contains all the vertices v' such that $p_{v'}$ passes through v. We set

$$A_v \;:=\; \{\; v' \mid (v, v') \in E(T) \;\}.$$

For each vertex v' there exists a unique vertex $v = \mathrm{pre}(v')$ such that $v' \in A_v$, called the *predecessor* of v'. A vertex $v \in V(T) - \{v_0\}$ will be called a *leaf* if $A_v = \emptyset$. We write $B(T)$ for the set of leaves of T.

Definition 2.5. An *order* on T is a bijection $\psi : \{1, \ldots, r\} \xrightarrow{\sim} B(T)$ such that for each $v \in V(T)$ the set

$$I_v \;:=\; \{\; i \;\mid\; \psi(i) \in V(T_v) \;\} \subset \{1, \ldots, r\}$$

is an interval, i.e. $I_v = \{r', \ldots, r''\}$. The triple (T, v_0, ψ) is called an *ordered tree.*

An order ψ on T induces a strict ordering of the sets A_v: for $v', v'' \in A_v$ we write $v' < v''$ if $i \in I_{v'}$ and $j \in I_{v''}$ implies $i < j$. Conversely, if we choose a strict ordering on each set A_v then there exists a unique order ψ on T inducing these orderings. In particular, an order ψ always exists. An automorphism of the ordered tree (T, v_0, ψ) is an automorphism $\kappa : T \xrightarrow{\sim} T$ such that $\kappa(v_0) = v_0$. Via ψ, κ induces a permutation of $\{1, \ldots, r\}$. Note that κ is uniquely determined by this permutation.

Let $\Pi = <\gamma_1, \ldots, \gamma_r \mid \prod \gamma_i = 1>$ be the free profinite group defined earlier, and let (T, v_0, ψ) be an ordered tree. For each $v \in V(T)$ we define an element

$$\gamma_v := \prod_{i \in I_v} \gamma_i$$

and a subgroup

$$\Pi_v := <\gamma_{v'} \mid v' \in A_v> \subset \Pi$$

of Π. Note that $\gamma_{v_0} = 1$, $\gamma_{\psi(i)} = \gamma_i$ and that for all $v \in V(T)$ we have

$$\gamma_v = \prod_{v' \in A_v} \gamma_{v'}, \tag{2.7}$$

where the product is taken with respect to the ordering on A_v induced by ψ.

Definition 2.6. An automorphism $\tau : \Pi \xrightarrow{\sim} \Pi$ is *compatible* with (T, v_0, ψ) if there exists an automorphism κ of (T, v_0, ψ), an element $\chi \in \hat{\mathbb{Z}}^\times$ and elements $\alpha_v \in \Pi_{\mathrm{pre}(v)}$, for all $v \in V(T)$, such that the following holds. For each $v \in V(T) - \{v_0\}$, let $\beta_v := \alpha_{v_1} \cdot \ldots \cdot \alpha_v$, where $p_v = (v_0, v_1, \ldots, v)$ is the shortest path from v_0 to v. Then

$$\tau(\gamma_v) = \beta_{\kappa(v)}\, \gamma_{\kappa(v)}^{\chi}\, \beta_{\kappa(v)}^{-1}.$$

It is easy to see that $\chi = \chi(\tau)$ and $\kappa = \kappa(\tau)$ are uniquely determined by τ.

Example 2.7. Let $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ be a presentation, as in Definition 1.4. An element $\sigma \in \Gamma_k$ induces an automorphism of $\pi_1(U)$, and hence, via ρ, an automorphism of Π. Let $V := \{0, 1, \ldots, r\}$, $E := \{(0,1), \ldots, (0,r)\}$ and $T := (V, E)$. We declare $0 \in V$ to be the root of T and let $\psi : \{1, \ldots, r\} \xrightarrow{\sim} B(T)$ be the identity. Then $(T, 0, \psi)$ is an ordered tree, and the automorphism $\Pi \xrightarrow{\sim} \Pi$ induced by $\sigma \in \Gamma_k$ is compatible with $(T, 0, \psi)$, by (1.3).

Definition 2.8. Let $T = (T, v_0, \psi)$ be an ordered tree. A *topological realization* of T consists of

- pairwise distinct points $z_0, \ldots, z_r \in \mathbb{P}^1_{\mathbb{C}}$ (we set $U := \mathbb{P}^1_{\mathbb{C}} - \{z_1, \ldots, z_r\}$),
- pairwise disjoint, connected, open subsets $U_v \subset U$, for $v \in V(T)$, and
- pairwise disjoint simple closed arcs $c_e : [0,1] \to U$, for $e \in E(T)$,

such that the following holds. The open subsets U_v are the connected components of $U - (\cup_e c_e)$. For each edge $e = (v_1, v_2)$, the arc c_e lies on the boundary of U_{v_1} and U_{v_2}, encircling U_{v_1} in clockwise and U_{v_2} in counterclockwise direction. For $i = 1, \ldots, r$, the set $U_{\psi(i)} \cup \{z_i\}$ is homeomorphic to an open disk. Finally, $z_0 \in U_{v_0}$.

Fix $v \in V(T) - v_0$, let $p_v = (v_0, v_1, \ldots, v_{k-1}, v)$ be the shortest path from v_0 to v, and set $e := (v_{k-1}, v)$. Set $z_v := c_e(0)$ and $z_{v_0} := z_0$. Choose a simple path $a_v : [0,1] \to \bar{U}_v$ leading from $z_{v_{k-1}}$ to z_v. Define

$$b_v \;:=\; a_{v_1} \cdots a_v, \quad \text{and} \quad \gamma_v \;:=\; b_v\, c_e\, b_v^{-1}.$$

We regard γ_v as an element of $\pi_1(U, z_0)$. To simplify the notation, set $\gamma_i := \gamma_{\psi(i)}$, for $i = 1, \ldots, r$, and $\gamma_{v_0} := 1$.

Lemma 2.9. *One can choose the paths a_v such that*

$$\gamma_v \;=\; \prod_{v' \in A_v} \gamma_{v'}$$

holds for all $v \in V(T)$ (in particular, $\gamma_1 \cdots \gamma_r = 1$) and that the elements $\gamma_1, \ldots, \gamma_r$ induce a presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$.

A collection (a_v) of paths as in Lemma 2.9 will be called a *skeleton* of the topological realization (z_i, U_v, c_e). The presentation ρ in Lemma 2.9 will be said to be induced by (a_v).

Proposition 2.10. *Let (z_i, U_v, c_e), (a_v) and ρ be as before. Let $\tau : U \xrightarrow{\sim} U$ be an orientation preserving diffeomorphism of U. Assume that $\tau(z_0) = z_0$ and that for all $v \in V(T)$ the restriction of τ to U_v is a diffeomorphism $U_v \xrightarrow{\sim} U_{v'}$, for some $v' \in V(T)$. Then the induced automorphism of $\pi_1(U, z_0) \cong \Pi$ is compatible with T.*

Proof. It is clear that τ induces an automorphism κ of T. We will use the notation $v' := \kappa(v)$. Replacing τ by a homotopic diffeomorphism, we may assume that $\tau(c_e) = c_{e'}$, for all $e \in E(T)$ (in particular, $\tau(z_v) = z_{v'}$). Fix $v \in V(T) - v_0$, let $p_v = (v_0, \dots, v_{k-1}, v)$ be the shortest path from v_0 to v and set $e := (v_{k-1}, v)$. Define

$$\alpha_{v'} \ := \ b_{v'_{k-1}} \, \tau(a_v) \, a_{v'}^{-1} \, b_{v'_{k-1}}^{-1}$$

and

$$\beta_{v'} \ := \ \tau(b_v) \, b_{v'}^{-1} = \alpha_{v'_1} \dots \alpha_{v'}$$

Since the closed path $\tau(a_v)\, a_{v'}$ lies entirely in $\bar{U}_{v'_{k-1}}$, $\alpha_{v'}$ is an element of the subgroup $\Pi_{v'_{k-1}}$. Moreover, we have

$$\tau(\gamma_v) \ = \ \tau(b_v)\, c_{v'}\, \tau(b_v)^{-1} \ = \ \beta_{v'}\, \gamma_{v'}\, \beta_{v'}^{-1}.$$

Therefore, τ is compatible with T (note that $\chi = 1$). □

2.3. The fundamental group of a degenerating curve. Let k be as in Section 2.1 and $S \subset \mathbb{P}^1_{k^s}$ a finite, Γ_k-invariant set and z_0 a k-rational base point for $U := \mathbb{P}^1_{k^s} - S$. In this section, we associate to (S, z_0) an ordered tree T. The tree T encodes the way in which the points in $S \cup \{z_0\}$ coalesce on the special fiber. If k_0 has characteristic 0 then there exists a presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ such that the Γ_k-action on $\pi_1(U, z_0) \cong \Pi$ is compatible with the tree T, in the sense of Definition 2.6 above. If k_0 is a subfield of $\mathbb{C}$, we can actually write down such a presentation. Moreover, we can compute the action of the inertia group on $\pi_1(U, z_0)$ explicitly, in terms of the *braid action.*

The valuation v of k extends uniquely to a valuation on k^s, which we also denote by v. Let R be the ring of integers of k^s. We identify the residue field of R with the algebraic closure k_0^s of k_0. Let $(X_R; z_{R,0}, \dots, z_{R,r})$ be the (unique) model over R of $(\mathbb{P}^1_{k^s}; z_0, \dots, z_r)$ as a *stable* $(r+1)$*-pointed tree of projective lines*, in the sense of [**12**]. We denote by $\bar{X} := X_R \otimes_R k_0^s$ the special fiber. By definition, $\bar{X}$ is a tree of projective lines, i.e. each irreducible component of $\bar{X}$ is non-canonically isomorphic to $\mathbb{P}^1$. For a point $z \in S \cup \{z_0\}$, we denote by $\bar{z}$ its specialization to $\bar{X}$. The points $\bar{z}_0, \dots, \bar{z}_r$ are pairwise distinct, smooth points of $\bar{X}$. Since the model X_R is unique, the natural Γ_k-action on $\mathbb{P}^1_{k^s}$ extends to an action on X_R. This yields an action of Γ_k on $\bar{X}$. In particular, the inertia group $I \lhd \Gamma_k$ acts k_0^s-linearly on $\bar{X}$.

We define the tree T as follows. The set of vertices $V(T)$ is the union of S with the set of irreducible components of $\bar{X}$. For every singular point x of $\bar{X}$, we draw an edge between the two vertices corresponding to the components meeting in x. Moreover, for each $z \in S$, we draw an edge between the vertex corresponding to z and the vertex corresponding to the component containing $\bar{z}$. We declare the component $\bar{X}_0$ which contains $\bar{z}_0$ as the root of T. It is clear that the elements of S are the leaves of T. If $v \in V(T)$ is not a leaf, we write $\bar{X}_v$ for the corresponding component of $\bar{X}$. The numbering $\{z_1, \dots, z_r\}$ of the set S corresponds to a bijection $\psi : \{1, \dots, r\} \xrightarrow{\sim} S = B(T)$. By changing this numbering, we may assume that ψ is an order. From now on, we fix ψ and regard $T = (T, \bar{X}_0, \psi)$ as an ordered tree. We remark that the Γ_k-action on $\bar{X}$ induces an action on T. It is clear that this action is determined by the action of Γ_k on $S = B(T)$.



Figure 2. The curve $\bar X$ and the associated tree T

Example 2.11. Let $k := \mathbb{Q}((t))$, $d \in \mathbb{Z}$, $S := \{\pm\sqrt{d} \pm i\sqrt{t}\}$ and $z_0 := 0$. The curve $\bar X$ consists of three components $\bar X_0$, $\bar X_5$ and $\bar X_6$. Each of these components corresponds to the choice of a coordinate w which identifies the function field of $\mathbb{P}^1_{k^s}$ with $k^s(w)$, modulo the action of $\mathrm{PGL}_2(R)$. For the component $\bar X_0$, we can choose the standard coordinate $w_0 := z$. For $\bar X_5$, we choose $w_5 := (z + \sqrt{d})/\sqrt{t}$, and for $\bar X_6$ we choose $w_6 := (z - \sqrt{d})/\sqrt{t}$. We let $z_1 := -\sqrt{d} + i\sqrt{t}$, $z_2 := -\sqrt{d} - i\sqrt{t}$, $z_3 := \sqrt{d} - i\sqrt{t}$ and $z_4 := \sqrt{d} + i\sqrt{t}$. Then z_1, z_2 reduce to $\bar z_1, \bar z_2 \in \bar X_5$ and z_3, z_4 reduce to $\bar z_3, \bar z_4 \in \bar X_6$. See Figure 2. We remark that the standard generator q_0 of the inertia group I (which sends $t^{1/n}$ to $e^{2\pi i/n}t^{1/n}$) acts as an involution on $\bar X_5$ and $\bar X_6$.

The following theorem can be thought of as a rigid-analytic analogue of Proposition 2.10. Many similar results can be found in the literature, see e.g. [**18**] or [**19**].

Theorem 2.12. *Assume that* $\mathrm{char}(k_0) = 0$. *There exists a presentation* $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ *such that the* Γ_k*-action on* Π *induced by* ρ *is compatible with* T.

Sketch of proof. We follow [**20**], with some modifications. The tree T can be equipped with a structure $\mathcal{G}$ of a *graph of groups*. For instance, to each vertex $v \in V(T)$ corresponding to a component $\bar X_v$ of $\bar X$ we associate the fundamental group $\pi_1(\bar U_v)$, where $\bar U_v$ is the open subset of $\bar X$ with all the points $\bar z_i$, $i > 0$, and all singular points removed. To each leaf $\psi(i) \in V(T)$, we associate the group $\hat{\mathbb{Z}}(1)$. We obtain a "canonical" isomorphism

$$\pi_1(U, z_0) \xrightarrow{\sim} \pi_1(T, \mathcal{G}). \tag{2.8}$$

In analogy to Lemma 2.9, one can choose a "skeleton" (a_v) of $(T, \mathcal{G})$ which induces a presentation $\Pi_r \xrightarrow{\sim} \pi_1(T, \mathcal{G})$ (actually, to make this precise, one has to consider $(T, \mathcal{G})$ as a *graphs of groupoids*). Let $\rho : \Pi_r \xrightarrow{\sim} \pi_1(U, z_0)$ be the composition of this presentation with the isomorphism (2.8). The rest of the proof of Theorem 2.12 is formally the same as the proof of Proposition 2.10. □

Remark 2.13. Assume that $\mathrm{char}(k_0) = p > 0$, and let $\pi_1^{\mathrm{adm}}(U, z_0)$ be the *admissible fundamental group*, i.e. the inverse limit over the finite quotients of $\pi_1(U, z_0)$ corresponding to Galois covers with admissible reduction (see [**20**]). There exists a surjective homomorphism $\rho : \Pi \to \pi_1^{\mathrm{adm}}(U, z_0)$ which induces an isomorphism on the maximal prime-to-p quotients such that the action of Γ_k on $\pi_1^{\mathrm{adm}}(U, z_0)$ is compatible with ρ and T, in an obvious sense. In particular, Theorem 2.12 remains true if we replace Π and $\pi_1(U, z_0)$ by their maximal prime-to-p quotients.

For a more detailed proof of Theorem 2.12 in a special case, see [**27**]. Under some extra assumptions, we can improve on Theorem 2.12 and define a concrete presentation ρ with the claimed properties. This presentation has the advantage that the inertia action on $\pi_1(U, z_0)$ is known explicitly.

We assume that k_0 is a subfield of the complex numbers, and that $k := k_0(t)^{\mathrm{h}}$ is the henselization of $k_0(t)$ at the place $t = 0$. For practical applications, this is not a serious restriction. We will regard k as a subfield of $\mathbb{C}((t))$ and its algebraic closure k^s as a subfield of $\mathbb{C}\{\{t\}\}$. We identify Γ_{k_0} as a subgroup of Γ_k, via its action on the coefficients of the Puiseux-expansion of elements of k. We let q_0 be the "canonical" generator of the inertia group $I \lhd \Gamma_k$, i.e. we have $q_0(t^{1/n}) = e^{2\pi i/n} t^{1/n}$. For simplicity, we assume moreover that $\infty \notin S \cup \{z_0\}$.

Let n be the smallest positive integer such that all points $z_0, \ldots, z_r$, regarded as elements of k^s, lie in $\mathbb{C}((t^{1/n}))$. We let $\tilde{t} := t^{1/n}$ and regard the z_i as germs of analytic functions of $\tilde{t}$. Choose $\epsilon > 0$ such that $z_0, \ldots, z_r$ are meromorphic on the disk

$$\tilde{D} \ := \ \{ \ \tilde{t} \in \mathbb{C} \mid \ |\tilde{t}| < \epsilon \ \}$$

and holomorphic on $\tilde{D}^* := \tilde{D} - \{0\}$. If ϵ is sufficiently small, the values $z_i(\tilde{t})$ are pairwise distinct, for all $\tilde{t} \in \tilde{D}^*$. The R-curve X_R gives rise to an analytic space $X_{\tilde{D}}$, together with a map $p : X_{\tilde{D}} \to \tilde{D}$ such that $p^{-1}(\tilde{D}^*) = \mathbb{P}^1_{\mathbb{C}} \times \tilde{D}^*$ and $p^{-1}(0) = \bar{X} \otimes \mathbb{C}$. Let $e = (v_1, v_2)$ be an edge corresponding to a singular point $x_e \in \bar{X}$. The complete local ring of x_e on X_R has the form

$$\hat{\mathcal{O}}_{X_R, x_e} \ = \ R[[\, u_e, v_e \ \mid \ u_e v_e = \tilde{t}^{n_e} \,]],$$

(for some well determined positive integer n_e) such that $u_e = 0$ (resp. $v_e = 0$) defines the component $\bar{X}_{v_1}$ (resp. $\bar{X}_{v_2}$) in a neighborhood of x_e. We may assume that u_e, v_e are analytic functions in a neighborhood of x_e on $X_{\tilde{D}}$. Choosing ϵ sufficiently small, we may identify the analytic set

$$V_e \ := \ \{ \ (u_e, v_e, \tilde{t}) \in \mathbb{C}^3 \ \mid \ u_e v_e = \tilde{t}^{n_e}, \ |u_e|, |v_e| < \epsilon^{n_e/2n} \ \}$$

with an open neighborhood of x_e on $X_{\tilde{D}}$. We may also assume that the sets V_e are pairwise disjoint.

Fix a positive real number t_0 such that $0 < t_0 < \epsilon$ and let $\tilde{z}_i := z_i(t_0^{1/n})$, for $i = 0, \ldots, r$, and

$$U_{t_0} \ := \ \mathbb{P}^1_{\mathbb{C}} - \{\tilde{z}_1, \ldots, \tilde{z}_r\} \ \subset \ p^{-1}(t_0^{1/n}).$$

For each edge e corresponding to a singular point x_e, define the closed arc

$$c_e \ := \ \left\{ \begin{array}{ccc} [0,1] & \longrightarrow & U_{t_0} \cap V_e \\ s & \mapsto & (t_0^{n_e/2n} e^{2\pi i s}, \, t_0^{n_e/2n} e^{-2\pi i s}, \, t_0^{1/n}) \end{array} \right. .$$

For the edge e adjacent to the leaf $\psi(i)$, let $c_e : [0,1] \to U_{t_0}$ be a small closed arc encircling $\tilde{z}_i$ in counterclockwise direction. We may assume that all the arcs c_e are pairwise disjoint. For $v \in V(T) - \{v_0\}$, set $e := (\mathrm{pre}(v), v) \in E(T)$ and let U_v be the connected component of $U_{t_0} - (\cup_{e'} c_{e'})$ containing the annulus $\{ \, (u_e, v_e, \tilde{t}) \in V_e \mid |u_e| > |v_e| \, \}$.

Lemma 2.14. *The data $(\tilde{z}_i, U_v, c_e)$ is a topological realization of the ordered tree T.*

Let us choose a skeleton (a_v) for $(\tilde{z}_i, U_v, c_e)$ and let $\rho_{t_0} : \Pi \xrightarrow{\sim} \pi_1(U_{t_0}, \tilde{z}_0)$ be the induced presentation (see Lemma 2.9). We define the presentation $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ as the composition of ρ_{t_0} with the canonical isomorphism

$$\pi_1(U_{t_0}, \tilde{z}_0) \xrightarrow{\sim} \pi_1(U, z_0). \tag{2.9}$$

A careful modification of the proof of Theorem 2.12 yields (compare with [**18**]):

Proposition 2.15. *Theorem 2.12 holds with the presentation ρ constructed above.*

Let $0 < |t| < \epsilon$ and choose a root $t^{1/n}$. The set

$$S_t := \{ z_1(t^{1/n}), \ldots, z_r(t^{1/n}) \} \subset \mathbb{P}^1_{\mathbb{C}}$$

depends only on t, since changing the root $t^{1/n}$ only permutes the points of S_t. Let $\theta : [0,1] \to \mathbb{C}$ be the closed arc $\theta(s) := t_0\, e^{2\pi i s}$. The map $s \mapsto S_{\theta(s)}$ corresponds to an element $Q \in \mathcal{B}_r$ of the *Artin braid group* on r strings. Here we identify $\mathcal{B}_r$ with the fundamental group of the space of r unordered points in U_{t_0}, with base point $S_{t_0} = \{\tilde{z}_1, \ldots, \tilde{z}_r\}$. There is a well known action of $\mathcal{B}_r$ on $\pi_1(U_{t_0}, \tilde{z}_0)$, see e.g. [**24**], Section II.10. By the definition of this action, the braid Q defined above corresponds, via the canonical isomorphism (2.9), to the inertia generator q_0 acting on $\pi_1(U, z_0)$. This gives us a practical way to compute this action explicitly, see [**9**].

Example 2.16. Let k, S, z_0 be as in Example 2.11. See Figure 3 for a picture of the presentation ρ_{t_0}, for $t_0 > 0$ sufficiently small and $d > 0$.

It is easily seen that the inertia generator q_0 corresponds to the braid $Q = Q_1 Q_3$ (where Q_1, Q_2, Q_3 are the standard generators of $\mathcal{B}_r$, see [**24**], Section II.10). Therefore, q_0 acts on Π via

$$^{q_0}\gamma_1 = \gamma_1\gamma_2\gamma_1^{-1}, \quad ^{q_0}\gamma_2 = \gamma_1, \quad ^{q_0}\gamma_3 = \gamma_3\gamma_4\gamma_3^{-1}, \quad ^{q_0}\gamma_4 = \gamma_3. \tag{2.10}$$

Set $\gamma_5 := \gamma_1\gamma_2$, $\gamma_6 := \gamma_3\gamma_4$, $\Pi_5 := <\gamma_1, \gamma_2> \subset \Pi$ and $\Pi_6 := <\gamma_3, \gamma_4> \subset \Pi$. Let $\sigma \in \Gamma_{\mathbb{Q}}$. By Theorem 2.12 and Proposition 2.15, the action of σ on $\Pi \cong \pi_1(U, z_0)$ is compatible with the tree T. We conclude that

$$^{\sigma}\gamma_i = \beta_{\sigma(i)}\, \gamma_{\sigma(i)}^{\chi(\sigma)}\, \beta_{\sigma(i)}^{-1}, \qquad \text{with} \quad \beta_i \in \begin{cases} \Pi_5, & \text{for } i = 1, 2 \\ \Pi_6, & \text{for } i = 3, 4. \end{cases} \tag{2.11}$$

Moreover, we have

$$^{\sigma}\gamma_5 = \gamma_5^{\epsilon(\sigma)\chi(\sigma)}, \qquad ^{\sigma}\gamma_6 = \gamma_6^{\epsilon(\sigma)\chi(\sigma)}, \tag{2.12}$$

where $\epsilon : \Gamma_{\mathbb{Q}} \to \{\pm 1\}$ is the Kummer character $\epsilon(\sigma) := {}^{\sigma}\sqrt{d}/\sqrt{d}$.

2.4. Computation of the residue. In this section we compute the residue $r_v(\omega)$ (and the value ω_0, if $r_v(\omega) = 0$) in a nontrivial example. Let us fix a square free integer d and let $k := \mathbb{Q}((t))$, $S := \{\pm\sqrt{d} \pm i\sqrt{t}\}$ and $z_0 := 0$, as in Example 2.11 and Example 2.16. Let $\rho : \Pi \xrightarrow{\sim} \pi_1(U, z_0)$ be the presentation constructed in the last subsection (see Figure 3). Then the action of Γ_k on $\pi_1(U, z_0) \cong \Pi$ is subject to (2.10), (2.11) and (2.12).



Figure 3.

Let $G := \tilde{A}_5$ be the nonsplit central extension of A_5 by $C := \{\pm 1\}$. We denote by $3A$ the (unique) conjugacy class of elements of order 3 in G and by $\mathrm{Ni}^{\mathrm{in}}(3A^4)$ the set of Nielsen classes $[\mathbf{g}] = [g_1, g_2, g_3, g_4]$ with $g_i \in 3A$, for $i = 1, \dots, 4$.

Proposition 2.17. *Given* $\mathbf{g} = (g_1, g_2, g_3, g_4)$ *such that* $[\mathbf{g}] \in \mathrm{Ni}^{\mathrm{in}}(3A^4)$*, let* $f : Y \to \mathbb{P}^1_{k^s}$ *be the* G*-cover with branch locus* S *and Hurwitz description* $\mathbf{g}$*, with respect to the presentation* ρ*. We denote by* k' *the smallest field containing* k *which is a field of moduli for* f *and by* $\omega \in H^2(k', C)$ *the f.o.d.-obstruction. Set* $g_5 := g_1 g_2 = (g_3 g_4)^{-1}$*. Then one of the three following cases occurs:*

- (i) *Case 1:* $g_5 = 1$*. We have* $k' = k = \mathbb{Q}((t))$ *and* $r_v(\omega) = (-d) \in \mathbb{Q}^*/\mathbb{Q}^{*2}$*. If* $d = -1$ *then* $r_v(\omega) = 0$ *and* $\omega_0 = (-1, -1) \in \mathrm{Br}_2(\mathbb{Q})$*.*
- (ii) *Case 2:* g_5 *has order* 10*. Then* $k' = \mathbb{Q}(\sqrt{5})((\tilde{t}))$*, with* $5v(\tilde{t}) = v(t)$*,* $r_v(\omega) = 0$ *and* $\omega_0 = (-1, -d) \in \mathrm{Br}_2(\mathbb{Q}(\sqrt{5}))$*.*
- (iii) *Case 3:* g_5 *has order* 6*. Then* $k' = \mathbb{Q}(\sqrt{-3})((\tilde{t}))$*, with* $3v(\tilde{t}) = v(t)$*,* $r_v(\omega) = 0$ *and* $\omega_0 = (-1, d) = (3, d) \in \mathrm{Br}_2(\mathbb{Q}(\sqrt{-3}))$*.*

Proof. There are exactly 18 classes $[g_1, g_2, g_3, g_4]$ in $\mathrm{Ni}^{\mathrm{in}}(3A^5)$. Among them, there are two classes with $g_1 g_2 = 1$, 10 classes with $g_5 := g_1 g_2$ of order 10 and 6 classes with g_5 of order 6. See e.g. [**1**, Prop. 5.8] for the use of a Theorem of Serre [**21**] showing how to compute the orders of products of odd order elements in $\tilde{A}_n$.

Case 1: $g_5 = 1$. The two Nielsen classes $[\mathbf{g}] \in \mathrm{Ni}^{\mathrm{in}}(3A^4)$ such that $g_1 g_2 = 1$ are permuted by an outer automorphism of G. Therefore, it suffices to consider one of them. So we assume that $\mathbf{g} = (g_1, g_1^{-1}, g_3, g_3^{-1})$ and that g_1 (resp. g_3) is the (unique) lift of $(1\,2\,3) \in A_5$ (resp. $(3\,4\,5) \in A_5$) to an element of order 3. By (2.10), we have

$$\mathbf{g}^{q_0} \;=\; (g_1^{-1}, g_1, g_3^{-1}, g_3) \;=\; a_0\, \mathbf{g}\, a_0^{-1}, \tag{2.13}$$

where $a_0 \in G$ is a lift of $(1\,2)(4\,5) \in A_5$. Note that a_0 is of order 4. Equation (2.11) implies that

$$\mathbf{g}^{\sigma} \;=\; \begin{cases} (\, g_1^{\chi_\sigma},\, g_1^{-\chi_\sigma},\, g_3^{\chi_\sigma},\, g_3^{-\chi_\sigma}\,), & \text{if } \epsilon_\sigma = 1 \\ (\, g_3^{\chi_\sigma},\, g_3^{-\chi_\sigma},\, g_1^{\chi_\sigma},\, g_1^{-\chi_\sigma}\,), & \text{if } \epsilon_\sigma = -1, \end{cases} \tag{2.14}$$

for all $\sigma \in \Gamma_{\mathbb{Q}}$ (we have used that the groups $G_5 := <g_1, g_1^{-1}>$ and $G_6 := <g_3, g_3^{-1}>$ are cyclic). It follows that $\mathbf{g}^{\sigma} = b_\sigma\, \mathbf{g}\, b_\sigma^{-1}$, where

$$b_\sigma \;:=\; a_0^{(1-\chi_\sigma)/2}\, b^{(1-\epsilon_\sigma)/2}, \tag{2.15}$$

and $b \in G$ is a lift of $(1\,4)\,(2\,5) \in A_5$. Together with (2.13), we get that $[\mathbf{g}]^{\tau} = [\mathbf{g}]$, for all $\tau \in \Gamma_k$. This shows that $k = \mathbb{Q}((t))$ is the field of moduli of the G-cover f.

In $\tilde{A}_5$, we have the equality $b\,a_0\,b^{-1} = a_0^{-1} = -a_0$. Using this and (2.15), we can compute the $\Gamma_{\mathbb{Q}}$-action on a_0 (defined by (2.2)):

$$\text{(2.16)} \qquad {}^{\sigma}a_0 \;:=\; b_\sigma\, a_0\, b_\sigma^{-1} \;=\; a_0^{\epsilon_\sigma}.$$

By Proposition 2.2 (i), the residue $r_v(\omega) \in H^1(\mathbb{Q}, \{\pm 1\})$ is represented by the cocycle

$$\text{(2.17)} \qquad c_\sigma \;=\; a_0\,({}^{\sigma}a_0)^{-\chi_\sigma^{-1}} \;=\; a_0^{1-\epsilon_\sigma\chi_\sigma^{-1}} \;=\; \epsilon_\sigma\eta_\sigma,$$

where $\eta_\sigma := {}^{\sigma}i/i$. Therefore, $r_v(\omega) = (-d) \in \mathbb{Q}^*/\mathbb{Q}^{*2} = H^1(\mathbb{Q}, \{\pm 1\})$. Since we assumed d to be square free, we have $r_v(\omega) = 0$ if and only if $d = -1$.

If $d = -1$, then $r_v(\omega) = 0$ and the value $\omega_0 \in H^2(\mathbb{Q}, \{\pm 1\}) = \mathrm{Br}_2(\mathbb{Q})$ of ω at v is well defined. By Proposition 2.2 (ii), ω_0 is the obstruction for lifting the homomorphism $\bar{\varphi} : \Gamma_{\mathbb{Q}} \to \bar{G} = A_5$ given by

$$\text{(2.18)} \qquad \bar{\varphi}(\sigma) \;=\; \bar{b}_\sigma \;=\; (1\,2)(4\,5)^{(1-\chi_\sigma)/2}\,(1\,4)(2\,5)^{(1-\epsilon_\sigma)/2}$$

to a homomorphism $\varphi : \Gamma_{\mathbb{Q}} \to \tilde{A}_5$. In other words, it is the obstruction for lifting the $\mathbb{Z}/2 \times \mathbb{Z}/2$-extension $\mathbb{Q}(\sqrt{3}, i)/\mathbb{Q}$ to a quaternion extension. As in the proof of Proposition 1.7, we conclude that

$$\text{(2.19)} \qquad \omega_0 \;=\; (-1,3) + (-1,-1) + (3,-1) \;=\; (-1,-1) \;\neq\; 0.$$

Case 2: g_5 has order 10. Let O be the set of Nielsen classes $[\mathbf{g}] \in \mathrm{Ni}^{\mathrm{in}}(3A^4)$ such that $g_5 := g_1 g_2$ has order 10. The inertia generator q_0, which acts on $\mathrm{Ni}^{\mathrm{in}}(3A^4)$, stabilizes O and has two orbits O_1 and O_2 of length 5, where a Nielsen class $[\mathbf{g}] \in O$ belongs to O_1 (resp. O_2) if $g_5 \in 10A$ (resp. $10B$). Here $10A$ and $10B$ are the two conjugacy classes in $\tilde{A}_5$ of elements of order 10. Since the classes $10A$ and $10B$ are conjugate over $\mathbb{Q}(\sqrt{5})$, it follows from (2.12) that $O_1^\sigma = O_2$ for $\sigma \in \Gamma_{\mathbb{Q}}$ with ${}^{\sigma}\sqrt{5} = -\sqrt{5}$. Therefore, Γ_k acts transitively on O, and the fixed field of any class $[\mathbf{g}] \in O$ is of the form $k' := \mathbb{Q}(\sqrt{5})((\tilde{t}))$, with $5v(\tilde{t}) = v(t)$. Hence, all classes $[\mathbf{g}]$ with g_5 of order 10 are conjugate under the action of Γ_k. We may therefore assume that $\mathbf{g} := (g_1, g_2, g_2^{-1}, g_1^{-1})$, where g_1 is a lift of $(1\,2\,3) \in A_5$ and g_2 is a lift of $(1\,4\,5)$. The G-cover f with Hurwitz description $[\mathbf{g}]$ has field of moduli k' as above. Therefore, there exists an element $a_0 \in G$ such that $\mathbf{g}^{q_0^5} = a_0\,\mathbf{g}\,a_0^{-1}$. In particular, $a_0\,g_5\,a_0^{-1} = g_5$. Since any subgroup of order 5 in A_5 is self-centralizing, the image of a_0 in A_5 has odd order. From Corollary 2.3 we conclude that $r_v(\omega) = 0$.

For all $\sigma \in \Gamma_{\mathbb{Q}(\sqrt{5})}$, there exists $b_\sigma \in G$ such that $\mathbf{g}^\sigma = b_\sigma\,\mathbf{g}\,b_\sigma^{-1}$. By (2.12), we have

$$\text{(2.20)} \qquad g_5^\sigma \;=\; g_5^{\chi_\sigma\epsilon_\sigma} \;=\; b_\sigma\, g_5\, b_\sigma^{-1}.$$

In particular, b_σ lies in N, the normalizer of $<g_5> \subset \tilde{A}_5$. Let $\bar{N}$ be the image of N in A_5. The group $\bar{N}$ is dihedral of order 10. Let $\bar{\varphi}(\sigma) := \bar{b}_\sigma$ and $\bar{\psi} : \Gamma_{\mathbb{Q}(\sqrt{5})} \to <\pm 1>$ the composition of $\bar{\varphi}$ with the sign character $\bar{N} \to \{\pm 1\}$. By (2.20), $\bar{\psi}(\sigma) \equiv \chi_\sigma\epsilon_\sigma \pmod 5$. A computation shows that $\bar{\psi}$ corresponds to the quadratic extension

$$\mathbb{Q}\big(\sqrt{5}, \sqrt{d(-10 + 2\sqrt{5})}\,\big) \;/\; \mathbb{Q}(\sqrt{5}).$$

To lift $\bar{\varphi}$ to $\tilde{A}_5$ it suffices to lift $\bar{\psi}$ to a homomorphism $\psi : \Gamma_{\mathbb{Q}(\sqrt{5})} \to \mathbb{Z}/4$ (compare with the proof of Proposition 1.7). Using $10 - 2\sqrt{5} = (1 - \sqrt{5})^2 + 2^2$, we conclude that

$$\omega_0 \;=\; \big(-1,\; d(-10+2\sqrt{5})\big) \;=\; (-1,-d).$$

Case 3: g_5 has order 6. Let $O \subset \mathrm{Ni}^{\mathrm{in}}(3A^4)$ be the set of classes $[\mathbf{g}]$ with g_5 of order 6. As in Case 2, q_0 acts on O and has two orbits, of length 3. Therefore, the fixed field of $[\mathbf{g}] \in O$ is of the form $k' = k_0((\tilde{t}))$, with $3v(\tilde{t}) = v(t)$ and $k_0/\mathbb{Q}$ at most a quadratic extension. We claim that $k_0 = \mathbb{Q}(\sqrt{-3})$, for all $[\mathbf{g}] \in O$. It suffices to show the following. Let $\sigma \in \Gamma_{\mathbb{Q}}$ such that $\mathbf{g}^\sigma = b_\sigma\, \mathbf{g}\, b_\sigma^{-1}$, for some $b_\sigma \in G$. Then $\chi(\sigma) \equiv 1 \pmod 6$.

The subgroups $G_5 := <g_1, g_2>$ and $G_6 := <g_3, g_4>$ of $\tilde{A}_5$ are of order 24, isomorphic to $\tilde{A}_4$ (two 3-cycles in A_5 generate either a cyclic subgroup of order 3 or a subgroup isomorphic to A_4). Moreover, $G_5 \cap G_6$ is cyclic of order 6, generated by g_5. Now assume that $\mathbf{g}^\sigma = b_\sigma\, \mathbf{g}\, b_\sigma^{-1}$ and $\epsilon(\sigma) = 1$. It follows from (2.11) that $b_\sigma\, G_5\, b_\sigma^{-1} = G_5$ and $b_\sigma\, G_6\, b_\sigma^{-1} = G_6$. It is easy to see that this implies $b_\sigma \in G_5 \cap G_6 = <g_5>$. Using (2.12), we get $g_5^{\chi(\sigma)} = b_\sigma\, g_5\, b_\sigma^{-1} = g_5$, hence $\chi(\sigma) \equiv 1 \pmod 6$. The argument is similar if $\epsilon(\sigma) = -1$. Then $b_\sigma\, G_5\, b_\sigma^{-1} = G_6$ and $b_\sigma\, G_6\, b_\sigma^{-1} = G_5$, and one can show that this implies $b_\sigma\, g_5\, b_\sigma^{-1} = g_5^{-1}$. Using again (2.12), we obtain $g_5^{-\chi(\sigma)} = b_\sigma\, g_5\, b_\sigma^{-1} = g_5^{-1}$, hence $\chi(\sigma) \equiv 1 \pmod 6$.

Let f be the G-cover with Hurwitz description $[\mathbf{g}] \in O$. We have shown that the minimal field of moduli of f containing k is of the form $k' = \mathbb{Q}(\sqrt{-3})((\tilde{t}))$, with $3v(\tilde{t}) = v(t)$. Therefore, $\mathbf{g}^{q_0^3} = a_0\, \mathbf{g}\, a_0$, for some $a_0 \in G$. As in Case 2, we conclude that $a_0 \in <g_5>$ and can therefore be assumed to be of odd order. By Corollary 2.3, we have $r_v(\omega) = 0$.

Recall that $\mathbf{g}^\sigma = b_\sigma\, \mathbf{g}\, b_\sigma$ implies $b_\sigma\, g_5\, b_\sigma^{-1} = g_5^{\epsilon(\sigma)}$. Therefore, the image of $\bar{\varphi}(\sigma) := \bar{b}_\sigma$ is contained in $\bar{N} \subset A_5$, the image of the normalizer of g_5 in $\tilde{A}_5$. The group $\bar{N}$ is dihedral of order 6, and the composition of $\bar{\varphi}$ with the character $\bar{N} \to \{\pm 1\}$ equals ϵ. As in Case 2, we conclude that $\omega_0 = (-1, d) = (3, d) \in \mathrm{Br}_2(\mathbb{Q}(\sqrt{-3}))$. □

2.5. Specialization to p-adic fields. In this section, we let k be a field which is complete with respect to a discrete valuation v, and assume that the residue field k_0 is perfect of characteristic $p > 0$. We denote by $\mathfrak{o}_k$ the ring of integers of k and by $\mathfrak{p}$ the maximal ideal of $\mathfrak{o}_k$. We set $K_t := k((t))$ and denote by v_t the valuation of K_t which has t as a uniformizer. We let $f_t : Y_t \to \mathbb{P}^1$ be a G-cover over $K_t^s = k^s\{\{t\}\}$ such that K_t is a field of moduli for f_t.

Our first goal is to define, for any element $a \in \mathfrak{p} - \{0\}$, the *specialization* $f_a : Y_a \to \mathbb{P}^1$ of f_t at $t = a$, which should be a G-cover over k^s with field of moduli k. Second, we would like to compute the residue $r_v(\omega_a)$ of the f.o.d.-obstruction ω_a of f_a in terms of a and the residue $r_{v_t}(\omega_t)$ of the f.o.d.-obstruction ω_t of f_t. Both these goals are problematic, in general, and we need some extra assumptions to succeed.

Let $A := \mathfrak{o}_k[[t]]$ and $K := \mathrm{Frac}(A)$. The ring A is a complete local domain, regular of dimension 2 and factorial. The field $K_t = k((t))$ is the completion of K at the valuation corresponding to the ideal $(t) \lhd A$. We denote this valuation also by v_t. Moreover, we identify K^s with the algebraic closure of K inside $K_t^s = k^s\{\{t\}\}$.

CONDITION 2.18.

(a) The branch locus $S_t \subset \mathbb{P}^1_{K_t^s}$ of the G-cover f_t is defined over K, i.e.

$$S_t \;=\; \{z \;\mid\; F(z) = 0\,\}, \qquad \text{with} \quad F(Z) \in K(Z).$$

(We may choose $F(Z) \in A(Z)$ such that the gcd of the coefficients is 1. We let $\delta(S) \lhd A$ be the ideal generated by the discriminant of F.)

(b) We have $\delta(S) = (t^n)$, with $n \geq 0$.

(c) The order of the group G is prime to $p = \mathrm{char}(k_0)$.

We assume from now on that Condition 2.18 is in force. By Condition 2.18 (a), the branch locus $S_t \subset \mathbb{P}^1_{K_s}$ of the G-cover f_t descends to a closed subscheme $S \subset \mathbb{P}^1_K$. Let $V := \mathrm{Spec}\, A[1/t]$ and $S_V \subset \mathbb{P}^1_V$ the closure of S inside $\mathbb{P}^1_V$. By Condition 2.18 (b), $S_V \to V$ is étale. Therefore, the embedding $S_V \subset \mathbb{P}^1_V$ corresponds to a morphism $V \to \mathcal{U}_r$, where $\mathcal{U}_r$ is the fine moduli space for the moduli problem "r distinct unordered points in $\mathbb{P}^1$". In more concrete terms, we have $\mathcal{U}_r = \mathbb{P}^r - \delta_r$ (δ_r denotes the discriminant hypersurface), and the coefficients of the polynomial $F(X) \in A[1/t][X]$ which defines S_V are the projective coordinates for the morphism $V \to \mathcal{U}_r$. Let $\mathcal{H}_r(G)$ be the Hurwitz space classifying G-Galois covers over $\mathbb{P}^1$ with exactly r branch points. To simplify the notation, we consider $\mathcal{U}_r$ and $\mathcal{H}_r(G)$ as schemes over $\mathfrak{o}$. Since the order of G is prime to the residue characteristic of $\mathfrak{o}$ (by Condition 2.18 (c)), the natural morphism $\mathcal{H}_r(G) \to \mathcal{U}_r$ (which associates to a G-cover its branch locus) is finite étale, see [**25**]. The G-cover f_t corresponds to a morphism $[f_t] : \mathrm{Spec}\, K_t \to \mathcal{H}_r(G)$, see Remark 1.3 (ii).

Lemma 2.19. *The morphism $V \to \mathcal{U}_r$ lifts to a morphism $\varphi : V \to \mathcal{H}_r(G)$ such that the following diagram commutes.*

$$\begin{tikzcd}
\mathrm{Spec}\, K_t \arrow[r, "{[f_t]}"] \arrow[d] & \mathcal{H}_r(G) \arrow[d] \\
V \arrow[r] \arrow[ur] & \mathcal{U}_r.
\end{tikzcd} \tag{2.21}$$

Proof. Since $\mathcal{H}_r(G) \to \mathcal{U}_r$ is a finite and étale morphism, the fiber product $V \times_{\mathcal{U}_r} \mathcal{H}_r(G)$ will be finite and étale over V. Let $V' \subset V \times_{\mathcal{U}_r} \mathcal{H}_r(G)$ be the irreducible component which is the image of the morphism $\mathrm{Spec}\, K_t \to V \times_{\mathcal{U}_r} \mathcal{H}_r(G)$. Then V' is integral, its fraction field K' is a finite extension of K and a subfield of K_t, and V' is the integral closure of V in K'. We have to show that $K' = K$.

Let A' be the integral closure of A in K'. Clearly, A' is a finite A-algebra, with fraction field K', and $A'[1/t]$ is étale over $A[1/t]$. Using Purity of Branch Locus and Abhyankar's Lemma, one shows that there exists an embedding $A' \hookrightarrow \mathfrak{o}'[[t^{1/n}]]$ of A-algebras, for some $n \geq 1$ and some finite unramified extension $\mathfrak{o}'/\mathfrak{o}$. In particular, the valuation v_t of K is totally ramified in K'. But K' was defined as a subfield of $K_t = k((t))$. Therefore, v_t is actually unramified in K'. Thus, by Purity of Branch locus, A'/A is étale. Since A is henselian, a finite étale extension of A is uniquely determined by its residue field extension. It follows that $A' = \mathfrak{o}'[[t]]$, for some finite unramified extension $\mathfrak{o}'/\mathfrak{o}$. Using again the embedding $A' \subset K_t = k((t))$, we conclude that $A' = A$. This proves the lemma. □

Lemma 2.19 shows in particular that f_t descends to a G-cover $f : Y \to \mathbb{P}^1_{K^s}$ over K^s such that K is a field of moduli for f, see Remark 1.3 (ii).

Let us fix an element a in $\mathfrak{p} - \{0\}$. We denote by v_a the discrete valuation of K corresponding to the ideal $(t - a) \lhd A$ (note that $v_a \neq v_t$). We choose an extension $\tilde{v}_a$ of v_a to K^s. Let $R \subset K^s$ denote the valuation ring corresponding to $\tilde{v}_a$. By Condition 2.18 (b), the branch locus $S \subset \mathbb{P}^1_{K^s}$ of f extends to a subscheme $S_R \subset \mathbb{P}^1_R$ such that S_R is étale over $\operatorname{Spec} R$. It follows that f has good reduction at $\tilde{v}_a$. Let $f_a : Y_a \to \mathbb{P}^1_{k^s}$ be the reduction of f at $\tilde{v}_a$. We call f_a the *specialization* of f_t at $t = a$. By definition, f_a is a G-cover defined over k^s, and k is a field of moduli for f_a.

The following theorem states that one obtains the f.o.d.-obstruction for f_a by 'specializing' the f.o.d.-obstruction for f_t at $t = a$. Moreover, this specialization procedure is compatible with computing the residue. The essential idea behind the theorem is that the class ω_t behaves just as one would expect it to behave if k were a local field of equal characteristic, with field of coefficients k_0, and the class ω_t arises from pull-back via the inclusion $k_0((t)) \hookrightarrow k((t))$.

Theorem 2.20. *Assume that Conditions 2.18 (a), (b) and (c) hold. Let $\omega_t \in H^2(K_t, C)$ be the f.o.d-obstruction for f_t and $\omega_a \in H^2(k, C)$ the f.o.d.-obstruction for f_a. Then*

(i) *The residue $r_t(\omega_t) \in H^1(k, C(-1))$ of ω_t at v_t is unramified at v, i.e. is induced by a class $\bar{r}_t(\omega_t) \in H^1(k_0, C(-1))$.*

(ii) *In $H^1(k_0, C(-1))$, we have the formula*

$$r_v(\omega_a) \;=\; v(a) \cdot \bar{r}_t(\omega_t).$$

(iii) *If $r_t(\omega_t) = 0$, then ω_t lies in the submodule $H^2(k_0, C)$, and the equality $\omega_a = \omega_t$ holds.*

Proof. Let $k^{\rm nr}$ be the maximal unramified extension of k and $\mathfrak{o}^{\rm nr}$ the ring of integers of $k^{\rm nr}$. We denote by $\mathcal{S}_A$ the set of discrete valuations of K which dominate A. Define $L^{\rm nr} \subset K^s$ as the maximal algebraic extension of K which is unramified over each valuation $w \in \mathcal{S}_A$. By Purity of Branch Locus, we can identify $\operatorname{Gal}(L^{\rm nr}/K)$ with $\pi_1(\operatorname{Spec} A)$, see [**13**]. Since A is a henselian local ring with residue field k_0, $\pi_1(\operatorname{Spec} A)$ is canonically isomorphic to Γ_{k_0}. Moreover, $B^{\rm nr} := \mathfrak{o}^{\rm nr}[[t]]$ is the integral closure of A in $L^{\rm nr}$. Similarly, let $L \subset K^s$ be the maximal algebraic extension of K which is unramified over each valuation $w \in \mathcal{S}_A - \{v_t\}$. We may identify $\operatorname{Gal}(L/K)$ with $\pi_1(V)$, where $V = \operatorname{Spec} A[1/t]$. Let B be the integral closure of A in L. Abhyankar's Lemma together with Purity shows that $B = \cup_n \mathfrak{o}^{\rm nr}[[t^{1/n}]]$, where n runs over all integers prime to p (compare with the proof of Lemma 2.19). Therefore, $\operatorname{Gal}(L^{\rm nr}/L) = \hat{\mathbb{Z}}'(1)$. We obtain the following commutative diagram:

$$\begin{tikzcd}
1 \arrow[r] & \hat{\mathbb{Z}}'(1) \arrow[r] \arrow[d] & \Gamma_{K_t} \arrow[r] \arrow[d] & \Gamma_k \arrow[r] \arrow[d] & 1 \\
1 \arrow[r] & \hat{\mathbb{Z}}'(1) \arrow[r] & \pi_1(V) \arrow[r] & \Gamma_{k_0} \arrow[r] & 1.
\end{tikzcd} \tag{2.22}$$

The vertical arrows are induced by the embedding of K^s into K_t^s. By [**13**], we have canonical isomorphism $H^i_{\rm \acute{e}t}(V, C) \cong H^i(\pi_1(V), C)$, for $i \geq 0$. Applying [**23**, II.A.1, §1 and §2] to (2.22), we obtain the following commutative diagram:

$$\begin{tikzcd}
0 \arrow[r] & H^2(k_0, C) \arrow[r] \arrow[d] & H^2_{\rm \acute{e}t}(V, C) \arrow[r, "\bar{r}_t"] \arrow[d] & H^1(k_0, C(-1)) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & H^2(k, C) \arrow[r] & H^2(K_t, C) \arrow[r, "r_t"] & H^1(k, C(-1)) \arrow[r] & 0.
\end{tikzcd} \tag{2.23}$$

The lower row is nothing else then the residue sequence (2.1) for the field K_t. The vertical arrows are the restriction homomorphisms from Galois cohomology.

Let $\varphi : V \to \mathcal{H}_r(G)$ be the morphism given by Lemma 2.19 and $\omega_V := \varphi^*\tilde{\omega} \in H^2_{\text{ét}}(V, C)$ the pull back of the global f.o.d.-obstruction $\tilde{\omega} \in H^2_{\text{ét}}(\mathcal{H}_r(G), C)$, see Remark 1.3 (iii) and [**25**]. By functoriality and (2.21), the morphism $H^2_{\text{ét}}(V, C) \to H^2(K_t, C)$ in (2.23) maps ω_V to ω_t, the f.o.d.-obstruction for f_t. Therefore, Part (i) of Theorem 2.20 follows from the commutativity of diagram (2.23).

Let K_a denote the completion of K with respect to the discrete valuation v_a and K_a^{nr} its maximal unramified extension. We may identify K_a with $k((t-a))$ and K_a^{nr} with $k^s((t-a))$. Moreover, we may identify $\text{Gal}(K_a^{\text{nr}}/K_a)$ with Γ_k, the absolute Galois group of k. Since v_a is unramified in the extension L/K, there exists an embedding $L \hookrightarrow K_a^{\text{nr}} = k^s((t-a))$. Explicitly, we choose a compatible system $a^{1/n} \in k^s$ of nth roots of $a \in k$ (where n runs over the integers prime to p), and define $L \hookrightarrow K_a^{\text{nr}}$ by

$$t^{1/n} \;\longmapsto\; a^{1/n}\,(1 + \frac{t-a}{a})^{1/n} \;=\; a^{1/n}\,(1 + \frac{t-a}{na} + \cdots). \tag{2.24}$$

Let $q_0 \in \hat{\mathbb{Z}}'(1)$ be a topological generator, corresponding to a compatible system (ζ_n) of nth roots of unity, where $p \not| n$. Let $\hat{\mathbb{Z}}'(1) \hookrightarrow \Gamma_k$ be the natural morphism; it sends q_0 to the automorphism of k which maps $\pi^{1/n}$ to $\zeta_n \pi^{1/n}$, where π is a uniformizer of k. Similarly, let $\hat{\mathbb{Z}}'(1) \hookrightarrow \pi_1(V) = \text{Gal}(L/K)$ be the morphism that sends q_0 to the automorphism of L which maps $t^{1/n}$ to $\zeta_n t^{1/n}$. It follows from (2.24) that the restriction homomorphism $\Gamma_k^t \to \text{Gal}(L/K) = \pi_1(V)$ induces a the diagram

$$\begin{tikzcd}
1 \arrow[r] & \hat{\mathbb{Z}}'(1) \arrow[r] \arrow[d, "\cdot v(a)"'] & \Gamma_k \arrow[r] \arrow[d] & \Gamma_{k_0} \arrow[r] \arrow[d] & 1 \\
1 \arrow[r] & \hat{\mathbb{Z}}'(1) \arrow[r] & \pi_1(V) \arrow[r] & \Gamma_{k_0} \arrow[r] & 1,
\end{tikzcd} \tag{2.25}$$

where the left vertical arrow sends q_0 to $q_0^{v(a)}$. The vertical arrow on the right is clearly the identity. We apply again [**23**, II.A.1, §1 and §2], this time to (2.25), and we obtain the following diagram.

$$\begin{tikzcd}
0 \arrow[r] & H^2(k_0, C) \arrow[r] \arrow[d] & H^2_{\text{ét}}(V, C) \arrow[r, "\bar{r}_t"] \arrow[d] & H^1(k_0, C(-1)) \arrow[r] \arrow[d, "\cdot v(a)"'] & 0 \\
0 \arrow[r] & H^2(k_0, C) \arrow[r] & H^2(k, C) \arrow[r, "r_v"] & H^1(k_0, C(-1)) \arrow[r] & 0.
\end{tikzcd} \tag{2.26}$$

The upper row is the same as in (2.23), the lower row is the residue sequence for the field k, see (2.1). The left and middle vertical arrows are the restriction homomorphism from Galois cohomology; in particular, the left vertical arrow is the identity. Checking the definition of the residue map (see the proof of Proposition 2.2), one finds that the vertical arrow on the right is multiplication by $v(a)$ and that the diagram (2.26) commutes. This proves Part (ii) and (iii) of the theorem. □

EXAMPLE 2.21. Let $f : Y \to \mathbb{P}^1$ be the $\tilde{A}_5$-covers with branch locus $S = \{\pm\sqrt{d} \pm i\sqrt{t}\}$ and Hurwitz description $[\mathbf{g}] = [g_1, g_1^{-1}, g_3, g_3^{-1}] \in \text{Ni}^{\text{in}}(3A^4)$ considered in Proposition 2.17, Case 1. We showed that $\mathbb{Q}((t))$ is a field of moduli for f and that $r_t(\omega) = (-d) \in \mathbb{Q}^*/\mathbb{Q}^{*2}$, where $\omega \in H^2(\mathbb{Q}((t)), C)$ is the f.o.d.-obstruction and r_t denotes the residue map corresponding to the valuation with parameter t.

Let $p > 5$ be a prime and assume that $p \not| d$. Choose an embedding of $\bar{\mathbb{Q}}$ into $\bar{\mathbb{Q}}_p$. We will consider f as a cover over $\bar{\mathbb{Q}}_p((t))$, with field of moduli $\mathbb{Q}_p((t))$. The discriminant of S is

$$\delta(S) \;=\; (\, 2^{12}(d^2 + 2dt + t^2)d^2t^2 \,) \;=\; (\, t^2 \,) \;\lhd\; \mathbb{Z}_p[[t]],$$

and Condition 2.18 holds. We may therefore specialize f at $t = p$, and we obtain an $\tilde{A}_5$-cover $f_p : Y_p \to \mathbb{P}^1$ over $\bar{\mathbb{Q}}_p$, with branch locus $S_p = \{\pm\sqrt{d} \pm \sqrt{p}\}$ and field of moduli $\mathbb{Q}_p$. Since $p \not| d$, the residue $r_t(\omega) = (-d) \in H^1(\mathbb{Q}_p, \pm 1)$ is unramified, as predicted by Part (i) of Theorem 2.20. By Part (ii), $r_p(\omega_p) = (-\bar{d}) \in H^1(\mathbb{F}_p, \pm 1) = \mathbb{F}_p^*/\mathbb{F}_p^{*2}$, where $\omega_p \in H^2(\mathbb{Q}_p, \pm 1) = \mathrm{Br}_2(\mathbb{Q}_p)$ is the f.o.d.-obstruction for f_p. In terms of the norm residue symbol, we have

$$\omega_p \;=\; (\, -d, -p \,)_p \;=\; \begin{cases} -1 & \text{if } -d \equiv x^2 \pmod{p}, \\ 1 & \text{if } -d \not\equiv x^2 \pmod{p}. \end{cases}$$

It is not hard to extend the computations of Section 2.4 to show that the f.o.d.-obstruction $\omega_t \in H^2(\mathbb{Q}((t)), \pm 1) = \mathrm{Br}_2(\mathbb{Q}((t))\,)$ is given by the Hilbert symbol $(-d, -t)$. Thus, to specialize ω_t to ω_p, one had to "plug in" $t = p$.

References

[1] P. Bailey and M. D. Fried, *Hurwitz monodromy, spin separation and higher levels of a modular tower*, this volume.

[2] S. Beckmann, *Ramified primes in the field of moduli of branched coverings of curves*, J. of Algebra **125** (1989), 236–255.

[3] K.S. Brown, *Cohomology of groups*, Graduate texts in Mathematics, no. 87, Springer-Verlag, 1994.

[4] K. Coombes and D. Harbater, *Hurwitz families and arithmetic Galois groups*, Duke Math. J. **52** (1985), no. 4, 821–839.

[5] P. Dèbes and J.-C. Douai, *Algebraic covers: Field of moduli versus field of definition*, Annales Sci. E.N.S. **30** (1997), 303–338.

[6] ———, *Local-global principles for algebraic covers*, Israel J. Math. **103** (1998), 237–257.

[7] P. Dèbes, J.-C. Douai, and M. Emsalem, *Familles de Hurwitz et cohomologie non abélienne*, Ann. Inst. Fourier **50** (2000), no. 1, 113–149.

[8] P. Dèbes and D. Harbater, *Fields of definition of p-adic covers*, J. reine angew. Math. **498** (1998), 223–236.

[9] M. Dettweiler, *Plane curve complements and curves on Hurwitz spaces*, Preprint, 2000.

[10] M. D. Fried and H. Völklein, *The inverse Galois problem and rational points on moduli spaces*, Math. Ann. **290** (1991), 771–800.

[Fri95a] M.D. Fried, *Introduction to Modular Towers: Generalizing the relation between dihedral groups and modular curves*, Proceedings AMS-NSF Summer Conference, vol. 186, 1995, Cont. Math series, Recent Developments in the Inverse Galois Problem, pp. 111–171.

[11] A. Fröhlich, *Orthogonal representations of Galois groups, Stiefel-Whitney classes and Hasse-Witt invariants*, J. Reine Angew. Math. **360** (1985), 84–123.

[12] L. Gerritzen, F. Herrlich, and M. van der Put, *Stable n-pointed trees of projective lines*, Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen **91** (1988), 131–163.

[13] A. Grothendieck, *Revêtement étales et groupe fondamental (SGA I)*, LNM, no. 224, Springer-Verlag, 1971.

[14] B. Huppert, *Endliche Gruppen I*, Grundlehren, no. 134, Springer-Verlag, 1967.

[15] Y. Ihara, *On the embedding of* $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ *into* $\widehat{GT}$, The Grothendieck Theory of Dessins d'Enfants (Leila Schneps, ed.), Lond. Math. Soc. Lecture Note Series, no. 200, Camb. Univ. Press, 1994, pp. 289–321.

[16] Y. Ihara and M. Matsumoto, *On Galois actions on profinite completions of braid groups*, Proceedings AMS-NSF Summer Conference, vol. 186, 1995, Cont. Math series, Recent Developments in the Inverse Galois Problem, 173–200.

[17] G. Malle and B. H. Matzat, *Inverse Galois theory*, Monographs in Mathematics, Springer, 1999.

[18] H. Nakamura, *Limits of Galois representations in fundamental groups along maximal degeneration of marked curves, I*, Amer. J. Math. **121** (1999), 315–358.

[19] F. Pop, $\frac{1}{2}$ *Riemann Existence Theorem with Galois action*, Algebra and Number Theory (G. Frey and J. Ritters, eds.), W. de Gruyter, Berlin, 1994, pp. 193–218.

[20] M. Saïdi, *Revêtements modérés et groupe fondamental de graphe de groupes*, Comp. Math. **107** (1997), 321–340.

[21] J.-P. Serre, *Relêvements dans* $\tilde{A}_n$, C. R. Acad. Sci. Paris **311** (1990), 477–482.

[22] J.-P. Serre, *Topics in Galois theory*, Research notes in mathematics, 1, Jones and Bartlett Publishers, 1992, Lecture notes prepared by Henri Darmon.

[23] ———, *Cohomologie Galoisienne*, 5. ed., LNM, no. 5, Springer-Verlag, 1994.

[24] H. Völklein, *Groups as Galois groups*, Cambridge Studies in Adv. Math., no. 53, Cambridge Univ. Press, 1996.

[25] S. Wewers, *Construction of Hurwitz spaces*, Thesis, Preprint No. 21 of the IEM, Essen, 1998.

[26] ———, *Deformation of tame admissible covers of curves*, Aspects of Galois Theory (H. Völklein, ed.), London Math. Soc. Lecture Note Series, no. 256, Cambridge Univ. Press, 1999, pp. 239–282.

[27] ———, *Rational boundary points on Hurwitz spaces*, preprint 1999, at http://www.math.upenn.edu/∼wewers.

University of Pennsylvania
E-mail address: `wewers@math.upenn.edu`